ANXIETY, DEPRESSION, AND RELATED CONDITIONS

All Classes (Updated 06/26/2024)

Row A on the following Anxiety, Depression, and Related Conditions Disposition Table allows only TWO of the following:

Anxiety

- Unspecified anxiety
- Generalized Anxiety Disorder
- Situational anxiety (also called adjustment disorder with anxiety)
- Social Anxiety Disorder

Depression

- Unspecified depression
- Postpartum depression
- Situational depression* (also called adjustment disorder with depressed mood)
 *For Situational Depression AME can also use the <u>Situational Depression Disposition</u> Table.
- Situational Anxiety and Depression (adjustment disorder with mixed anxiety and depressed mood)

Oth<u>er</u>

- Obsessive Compulsive disorder (OCD)
- Post Traumatic Stress Disorder (PTSD)
- V code^{*} (DSM)/Z code (ICD-10) table items

Note^{*}: V code (DSM)/Z code (ICD-10) table includes but is not limited to:

- Uncomplicated Bereavement;
- Relationship distress with spouse or intimate partner;
- Parent-Child relational problem; and/or
- Phase of Life Problem (problems adjusting to major life transitions such as newly retired, getting married, empty-nest, new career, becoming a parent, etc.

Review ALL Row A criteria including psychotherapy (past and current) and medication history.

For conditions not listed above, see Row B.

ANXIETY, DEPRESSION, AND RELATED CONDITIONS

All Classes (Updated 06/26/2024)

DISEASE/CONDITION	EVALUATION DATA	DISPOSITION
A. Up to TWO listed conditions treated with any combination of:	The AME should gather information regarding the diagnosis, severity, treatment, symptoms, and address ALL of the questions on the Anxiety,	ISSUE
Psychotherapy (current or historical)	Depression, and Related conditions Decision Tool for the AME.	Summarize this history, diagnosis, and annotate Block 60:
Medication: A single mental health medication last taken, prescribed, or recommended two (2) or more years ago. This includes when the treating physician changed medications	Note: For individuals who were deferred, but now may possibly meet all Row A criteria, use the Fast Track Pathway Decision Tool for Currently Deferred Cases. If all items on the decision tool are in the clear, "NO column, the AME may:	"Discussed the history of [insert diagnosis here], no positives to screening questions, and no concerns."
for better treatment response, provided only ONE medication	DO NOT USE THIS ROW IF (go to Row B):	If any " YES " answers
was taken at a time and last use was two	⊗ Previously on an SI/SC for SSRI use;	OR
or more years ago. ACCEPTABLE DIAGNOSES:	Recurrent episodes or symptoms. If the individual had multiple failed attempts to discontinue or stay off medication(s) over any period;	Any AME concerns: DEFER
Uncomplicated:	⊗ If the individual has any history of	
Anxiety: Generalized Anxiety Disorder	requiring more than ONE mental health medication at a time (e.g., dual or triple therapy); and/or	
 Situational anxiety (aka adjustment disorder with anxiety) Social Anxiety Disorder Unspecified anxiety 	Diagnosed with Major Depressive Disorder (MDD) or Persistent Depressive Disorder (dysthymia).	
Postpartum depression Situational depression* (aka adjustment disorder with depressed mood) Situational Anxiety and Depression (adjustment disorder with mixed anxiety and depressed mood) Unspecified depression	 NOTE^: V code (DSM)/Z code (ICD-10) table includes but is not limited to: Uncomplicated Bereavement; Relationship distress with spouse or intimate partner; Parent-Child relational problem; and/or Phase of Life Problem (problems adjusting to major life transitions such as newly retired, getting married, empty-nest, new career, becoming a parent, etc. 	

DISEASE/CONDITION	EVALUATION DATA	DISPOSITION
3. Other: Obsessive Compulsive disorder (OCD) Post-Traumatic Stress Disorder (PTSD) V code^ (DSM)/Z code (ICD-10) table items Note*: For Situational Depression - AME can use this table or the Situational Depression Disposition Table.		
B. All Others	Submit a current, detailed Clinical Progress Note	
Previously on an SI/SC for SSRI use; OR	generated from a clinic visit with the treating physician or specialist no more than 90 days before the AME exam.	DEFER Annotate (elements or findings) in
Recurrent episodes or symptoms. If the individual had multiple failed attempts to discontinue or stay off medication(s) over any period;		Block 60.
OR		
If the individual has any history of requiring more than ONE mental health medication at a time (e.g., dual or triple therapy);		
OR		
Diagnosed with Major Depressive Disorder (MDD) or Persistent Depressive Disorder (dysthymia);		
AND/OR		
Seasonal Affective Disorder		