

**DRAFT**

**Environmental Assessment**

## **BLUE ORIGIN NEW GLENN CADENCE INCREASE AT SPACE LAUNCH COMPLEX 36**

**CAPE CANAVERAL SPACE FORCE STATION, FLORIDA**

**Appendix A.1.2, Part 1**

**September 2026**

**Federal Aviation Administration  
Department of the Air Force  
United States Coast Guard  
National Aeronautics and Space Administration**

Unique ID: EAXX-021-12-14Y-1773227485



**Federal Aviation  
Administration**

*Photo credit Blue Origin*

**LIST OF APPENDICES**

Appendix A      Regulatory Consultations ..... A-1

# Acronyms and Abbreviations

---

| <b>Acronym</b>     | <b>Definition</b>                              |
|--------------------|------------------------------------------------|
| §/§§               | section/sections                               |
| ACHP               | Advisory Council on Historic Preservation      |
| AHA                | Aircraft Hazard Area                           |
| APE                | Area of Potential Effects                      |
| ASEL               | A-weighted sound exposure level                |
| BCA                | Biological and Conference Assessment           |
| BGEPA              | Bald and Golden Eagle Protection Act           |
| Blue Origin        | Blue Origin, LLC                               |
| CCP                | Comprehensive Conservation Plan                |
| CCSFS              | Cape Canaveral Space Force Station             |
| CDNL               | C-weighted day-night average sound level       |
| CFR                | Code of Federal Regulations                    |
| CH <sub>4</sub>    | methane                                        |
| CMP                | Coastal Management Program                     |
| CO <sub>2</sub>    | carbon dioxide                                 |
| CO <sub>2</sub> e  | carbon dioxide equivalent                      |
| COMDTINST          | Commandant Instruction                         |
| CZMA               | Coastal Zone Management Act                    |
| DAF                | Department of the Air Force                    |
| dB                 | decibel(s)                                     |
| dba                | A-weighted decibels                            |
| DNL                | day-night average sound level                  |
| DoD                | Department of Defense                          |
| DOT                | Department of Transportation                   |
| DoW                | Department of War                              |
| EO                 | Executive Order                                |
| EA                 | Environmental Assessment                       |
| EFH                | essential fish habitat                         |
| EIS                | Environmental Impact Statement                 |
| ERD                | Environmental Resources Document               |
| ERP                | Environmental Resource Permit                  |
| ESA                | Endangered Species Act                         |
| FAA                | Federal Aviation Administration                |
| FDEP               | Florida Department of Environmental Protection |
| FONSI              | Finding of No Significant Impact               |
| HAPC               | habitat areas of particular concern            |
| ILV                | integrated launch vehicle                      |
| INRMP              | Integrated Natural Resources Management Plan   |
| IPaC               | Information for Planning and Consultation      |
| KSC                | Kennedy Space Center                           |
| LAA                | Limited Access Area                            |
| L <sub>A,max</sub> | maximum A-weighted sound level                 |
| lbf                | million pounds-force                           |
| LC                 | Launch Complex                                 |
| LH <sub>2</sub>    | liquid hydrogen                                |
| L <sub>max</sub>   | maximum sound level                            |
| LNG                | liquefied natural gas                          |
| LOX                | liquid oxygen                                  |

| <b>Acronym</b>    | <b>Definition</b>                                                    |
|-------------------|----------------------------------------------------------------------|
| MBTA              | Migratory Bird Treaty Act                                            |
| MG                | million gallons                                                      |
| MGD               | million gallons per day                                              |
| MINWR             | Merritt Island National Wildlife Refuge                              |
| MMPA              | Marine Mammal Protection Act                                         |
| MSA               | Magnuson-Stevens Fishery Conservation and Management Act             |
| MT/yr             | metric tons per year                                                 |
| N <sub>2</sub> O  | nitrous oxide                                                        |
| NAAQS             | National Ambient Air Quality Standards                               |
| NAS               | National Airspace System                                             |
| NASA              | National Aeronautics and Space Administration                        |
| NEI               | National Emissions Inventory                                         |
| NEPA              | National Environmental Policy Act                                    |
| NHPA              | National Historic Preservation Act                                   |
| NMFS              | National Marine Fisheries Service                                    |
| No.               | Number                                                               |
| NOTAM             | Notice to Airmen                                                     |
| NOTMAR            | Notice to Mariners                                                   |
| NO <sub>x</sub>   | nitrogen oxides                                                      |
| NPDES             | National Pollutant Discharge Elimination System                      |
| NRHP              | National Register of Historic Places                                 |
| OSHA              | Occupational Safety and Health Administration                        |
| PDC               | project design criteria                                              |
| PLOC              | Programmatic Letter of Concurrence                                   |
| PM                | particulate matter                                                   |
| PM <sub>10</sub>  | particulate matter with a diameter equal to or less than 10 microns  |
| PM <sub>2.5</sub> | particulate matter with a diameter equal to or less than 2.5 microns |
| PSD               | Prevention of Significant Deterioration                              |
| psf               | pound per square foot                                                |
| RNA               | Regulated Navigation Area                                            |
| SCCC              | Spaceport Command and Control Center                                 |
| SHPO              | State Historic Preservation Officer                                  |
| SJRWMD            | St. Johns River Water Management District                            |
| SLC               | Space Launch Complex                                                 |
| SLD 45            | Space Launch Delta 45                                                |
| SpaceX            | Space Exploration Technologies Corp.                                 |
| U.S.              | United States                                                        |
| U.S.C.            | United States Code                                                   |
| USCG              | United States Coast Guard                                            |
| USEPA             | United States Environmental Protection Agency                        |
| USFWS             | United States Fish and Wildlife Service                              |
| USSF              | United States Space Force                                            |
| VOL               | vehicle operator license                                             |

## **Appendix A      Regulatory Consultations**

### **A.1      Endangered Species Act Section 7 Consultation**

## A.1.2 National Marine Fisheries Service Programmatic Letter of Concurrence



UNITED STATES DEPARTMENT OF COMMERCE  
National Oceanic and Atmospheric Administration  
NATIONAL MARINE FISHERIES SERVICE  
1315 East-West Highway  
Silver Spring, Maryland 20910

5/4/26

Refer to NMFS No: OPR-2025-02196

Ms. Stacey Zee  
Manager, Operations Support Branch  
U.S. Dept. Transportation, Federal Aviation Administration  
Office of Commercial Space Transportation  
800 Independence Ave SW, Suite 325  
Washington, DC 20591

RE: Reinitiation of the Programmatic Letter of Concurrence for Space Launch and Reentry Operations

Dear Ms. Zee:

On July 29, 2025, the National Marine Fisheries Service (NMFS) received the Federal Aviation Administration's (FAA) request, in partnership with U.S. Space Force (USSF) and National Aeronautics and Space Administration (NASA), to reinitiate the programmatic consultation (hereafter referred to as the programmatic). FAA, USSF, and NASA proposed changes to the programmatic, to better meet the changing and growing demands of the commercial space industry.

This response to your request was prepared by NMFS pursuant to section 7(a)(2) of the Endangered Species Act of 1973, as amended (ESA; 16 U.S.C. 1531 et seq.), implementing regulations at 50 CFR part 402, and agency guidance for preparation of letters of concurrence. On March 30, 2026, in *Center for Biological Diversity v. Burgum*, No. 24-cv-04651 (N.D. Cal.), the U.S. District Court for the Northern District of California vacated aspects of four provisions from the 50 CFR part 402 regulations governing interagency consultation under section 7 of the Endangered Species Act and reinstated the provisions that were previously in effect. Consistent with the Court's ruling, these are the governing provisions for this consultation:

- "*Effects of the action* refers to the direct and indirect effects of an action on the species or critical habitat, together with the effects of other activities that are interrelated or interdependent with that action, that will be added to the environmental baseline. The environmental baseline includes the past and present impacts of all Federal, State, or private actions and other human activities in the action area, the anticipated impacts of all proposed Federal projects in the action area that have already undergone formal or early section 7 consultation, and the impact of State or private actions which are contemporaneous with the consultation in process. Indirect effects are those that are caused by the proposed action and are later in time, but still are reasonably certain to occur. Interrelated actions are those that are part of a larger action and depend on the larger action for their justification. Interdependent actions are those that have no

independent utility apart from the action under consideration.”<sup>1</sup> (50 CFR §402.02 (2018)).

- 50 CFR §402.16(a): “(a) Reinitiation of consultation is required and shall be requested by the Federal agency or by the Service, where discretionary Federal involvement or control over the action has been retained or is authorized by law and . . .” (50 CFR §402.16(a) (2023)).

Some of the information and analyses considered in this document were prepared under the version of the 50 CFR part 402 regulations in effect prior to the Court’s order. We have reviewed the information and analyses and confirm that the content of that information and those analyses, including the identification and analyses of effects, would be the same under either version of the applicable regulations.

An ESA section 7(a)(2) programmatic consultation is a consultation addressing an agency’s multiple similar, frequently occurring or routine actions in particular geographic areas; or under a proposed program, plan, policy, or regulation that provides a framework for future proposed actions (50 CFR §402.02). Programmatic consultations assess the program bounds or limits, referred to as the framework of the program, to insure the present and future actions conducted within that framework are not likely to jeopardize the continued existence of listed species or result in the destruction or adverse modification of critical habitat. This is a programmatic consultation on an informal programmatic action.

This consultation is a reinitiation of the 2023 Amended Programmatic Concurrence Letter for Launch and Reentry Vehicle Operations in the Marine Environment and Starship-Super Heavy Launch Vehicle Operations at SpaceX’s Boca Chica Launch Site, Cameron County, TX (OPR-2021-02908). Section 7(a)(2) of the ESA requires action agencies to consult with the Services, which is a continuing obligation for as long as the Federal action agency has discretion over the action and listed species or critical habitat may be affected. The reinitiation triggers for informal consultations are: 1) new information reveals effects of the action that may affect an ESA-listed species or designated critical habitat in a manner or to an extent not previously considered; 2) the identified action is subsequently modified in a manner that causes an effect to the ESA-listed species or designated critical habitat that was not considered in the concurrence letter; or if 3) a new species is listed or critical habitat designated that may be affected by the identified action. In those situations, where the action is ongoing, and the Federal action agency retains discretionary involvement or control over the action, it is appropriate to reinitiate consultation to satisfy the action agency’s section 7(a)(2) responsibilities. This consultation is being reinitiated because the action has been subsequently modified in a manner that causes an effect to ESA-listed species or critical habitat that was not considered in the previous programmatic concurrence letter, and because new critical habitat was designated that may be affected by the action.

---

<sup>1</sup> This definition includes the second sentence of the definition of “effects of the action.” That sentence provided the definition of “environmental baseline” in effect as of 2018. In the 2019 rule amending the 50 CFR part 402 regulations, the Services established “environmental baseline” as a stand alone definition. 84 Fed. Reg. 44976, 45016 (August 27, 2019). The Court’s ruling did not touch upon that definition of “environmental baseline,” and it therefore remains valid. The definition is also fully consistent with the definition of “effects of the action” from 2018.

The following species considered in this consultation are regulated under the Marine Mammal Protection Act (MMPA) and the ESA: the beluga whale (*Delphinapterus leucas*) – Cook Inlet Distinct Population Segment (DPS), blue whale (*Balaenoptera musculus*), bowhead whale (*Balaena mysticetus*), fin whale (*Balaenoptera physalus*), gray whale (*Eschrichtius robustus*) – Western North Pacific DPS, humpback whale (*Megaptera novaeangliae*) – Cape Verde Islands/Northwest Africa DPS, Central America DPS, Mexico DPS, and Western North Pacific DPS, killer whale (*Orcinus orca*) – Southern Resident DPS, North Atlantic right whale (*Eubalaena glacialis*), North Pacific right whale (*Eubalaena japonica*), rice's whale (*Balaenoptera ricei*), sei whale (*Balaenoptera borealis*), Southern right whale (*Eubalaena australis*), sperm whale (*Physeter macrocephalus*), bearded seal (*Erignathus barbatus nauticus*) – Beringia DPS, Guadalupe fur seal (*Arctocephalus townsendi*), ringed seal (*Phoca hispida hispida*) – Arctic subspecies, and Steller sea lion (*Eumetopias jubatus*) – Western DPS. Each statute defines take distinctly. The MMPA defines take as to harass, hunt, capture, or kill, or attempt to harass, hunt, capture, or kill any marine mammal (16 U.S.C. §1632(13)). Take under the ESA is to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any such conduct (16 U.S.C. §1532(19)). Actions considered 'take' under one statute do not necessarily rise to the level of take under the other statute.

This programmatic letter of concurrence underwent pre-dissemination review using standards for utility, integrity, and objectivity in compliance with applicable guidelines issued under the Data Quality Act (DQA; section 515 of the Treasury and General Government Appropriations Act for Fiscal Year 2001, Public Law 106-554). A complete record of this consultation is on file electronically with the NMFS Office of Protected Resources in Silver Spring, Maryland.

## CONSULTATION HISTORY

The NMFS provided a programmatic letter of concurrence (PLOC) to the FAA, USSF, and NASA for launch and reentry vehicle operations and Space Exploration Technologies Corporation (SpaceX) Starship-Super Heavy operations in the marine environment on January 31, 2022, and subsequently amended it on April 14, 2023 to clarify information and to include a Bahamas and Caribbean area in the PLOC. On August 6, 2024, the PLOC was amended to include Morro Bay Harbor, and associated conservation measures, as a port for commercial space vessel operations. On September 11, 2024, the PLOC was amended to add waters within a 90-mile (mi) or 145-kilometer (km) radius around Bermuda to the action area. On January 17, 2025, NMFS provided a conference and biological opinion to FAA for SpaceX Starship-Super Heavy operations, which superseded all Starship-Super Heavy operations considered in the PLOC. Therefore, Starship-Super Heavy operations are not covered under this programmatic. On June 25, 2025, the PLOC was amended to increase the maximum number of annual spacecraft reentry and landing operations in the Pacific Ocean portion of the action area from three to 11. Following are notable dates in the development of this latest reinitiation of the PLOC:

- **May 28, 2025:** FAA, as the lead action agency, presented the action agencies' proposed changes for the programmatic to NMFS to expand the current scope of the PLOC. Proposed changes included expanding the action area to the whole globe, updating launch and reentry vehicle specifications envelopes, and increasing the number of operations.

- **June 5–July 18, 2025:** NMFS and FAA engaged in pre-consultation and technical assistance.
- **July 29, 2025:** FAA formally requested reinitiation of the programmatic via email to NMFS.
- **August 21–September 12, 2025:** NMFS and FAA met multiple times to discuss timing considerations for the programmatic, consistent with Executive Order 14335, “Enabling Competition in the Commercial Space Industry,” and the activities that should be covered under the programmatic.
- **November 26, 2025:** NMFS requested, via email to FAA, additional information to analyze the action agencies’ proposed changes to the programmatic. On December 3, 2025, NMFS and FAA met to discuss the information.
- **January 29, 2026:** NMFS presented an analysis of the action agencies’ proposed changes to FAA.
- **February 19, 2026:** NMFS sent another request to FAA for additional information needed to initiate consultation. NMFS and FAA met to discuss the outstanding items needed for initiation. On February 20, 2026, FAA provided responses to our December 2025 and February 2026 requests for additional information via email.
- **February 23, 2026:** As requested, NMFS emailed the Proposed Action section of the programmatic to the FAA, with requests for additional information and for FAA’s review to ensure accuracy in the description of the proposed activities.
- **March 2, 2026:** FAA provided their review of the Proposed Action and responses to our requests for additional information via email. On March 3, 2026, NMFS and FAA met to discuss the proposed action and how the programmatic will supersede other space consultations.
- **March 4–March 11, 2026:** FAA and NMFS met multiple times to discuss adjustments to the proposed action, including the consultations to be superseded and an expansion of the action area. FAA also provided additional information related to those adjustments.

Previous ESA section 7 consultations for the activities that are included in and superseded by this reinitiated programmatic consultation include:

- **OPR-2025-02194:** On July 29, 2025, NMFS received a request for consultation from FAA for the proposed licensing of testing activities of Blue Origin’s launch vehicle, New Glenn. The proposed activities consisted of 1) New Glenn first stage testing, and 2) experiments on New Glenn’s payload fairings. First stage testing activities will include one explosive event off North Carolina and three explosive events off Florida. Fairing experiments will include expending some and recovering some of the fairings and instruments housed in the fairings. NMFS issued a letter of concurrence for the proposed activities on September 29, 2025.
- **OPR-2025-01425:** On May 15, 2025, NMFS received FAA’s request for concurrence on proposed Rocket Lab licensing activities, consisting of up to three launches of Rocket Lab’s Neutron launch vehicle between 2025 and 2026, including controlled first stage landing and disposal in the North Atlantic Ocean and second stage reentry, landing, and debris in the southern Indian Ocean and Southern Ocean. On July 14, 2025, NMFS issued an expedited letter of concurrence for those proposed activities.
- **OPR-2024-02184:** On August 27, 2024, NMFS received FAA’s request for concurrence for proposed Sierra Space Corporation (Sierra Space) licensing activities, consisting of

- four Dream Chaser (Sierra Space's spacecraft) reentries per year 2025–2029. Each reentry included expending Dream Chaser's cargo module in the Pacific Ocean. On June 6, 2025, NMFS issued a letter of concurrence for those proposed activities.
- **OPR-2024-01961:** On August 14, 2024, NMFS received FAA's request for concurrence on proposed Boeing Company (Boeing) licensing activities, consisting of up to four Commercial Space Transport 100, also known as Starliner (Boeing's spacecraft), reentries per year 2024–2028. Each reentry included expending Starliner's service module in the Pacific Ocean. On November 14, 2024, NMFS issued a letter of concurrence for those proposed activities.
  - **OPR-2024-01272:** On June 10, 2024, NMFS received FAA's request for concurrence on proposed Varda Space Industries, Inc. (Varda) licensing activities, consisting of three reentry, landing, and recovery operations of Varda's Winnebago series capsules, including expending of a Rocket Lab Corporation (Rocket Lab) satellite bus ('Pioneer'), in the Great Australian Bight. On August 20, 2024, NMFS issued a letter of concurrence for those proposed activities. On February 9, 2026, FAA relayed that Varda reentry, landing, and recovery operations will increase from three to 10. Given the small increase in operations in the same action area, NMFS concludes that this will not alter the effects analysis completed in OPR-2024-01272.
  - **WCRO-2024-00812:** On March 21, 2024, NMFS West Coast Regional Office received a request for consultation from USSF on an annual cadence of 100 launches per year of SpaceX Falcon 9 and Falcon Heavy from VSFB, 36 first stage recoveries per year, and 10 expended Falcon 9 or Falcon Heavy center core boosters per year. The action area was also expanded for dronship landings and fairing recovery operations. An expedited letter of concurrence was issued on April 17, 2024.
  - **WCRO-2023-00002:** On December 19, 2022, NMFS West Coast Regional Office received a request for consultation from USSF to increase the annual cadence of SpaceX Falcon 9 operations to 110 launches per year from VSFB and 36 first stage recoveries per year, with an expanded area for dronship landings and fairing recovery operations. An expedited letter of concurrence was issued on January 20, 2023.

## PROPOSED ACTION AND ACTION AREA

When consulting on a national program, agencies are often not able to provide enough specificity about the number, location, timing, frequency, precise methods, and intensity of each action's activities to determine the site-specific effects those activities will have on ESA-listed species and critical habitat. This letter of concurrence documenting the programmatic consultation includes a review of the effects of activities covered by the FAA, USSF, and NASA space launch and reentry actions and establishes a framework for authorizing future actions and tracking their effects.

This programmatic consultation is multifaceted and consists of activities ranging from launch, reentry, and landing operations, pre- and post-launch surveillance, and recovery operations. The following sections provide an overview of the action agencies and their authorities pertaining to this consultation, the activities covered by this consultation, and conservation measures FAA, USSF, and NASA will implement to minimize the effects of the space launch and reentry program on ESA-listed or proposed species and designated or proposed critical habitat.

## Action Agency Overview

### *Federal Aviation Administration*

The FAA Office of Commercial Space Transportation oversees, licenses, and regulates U.S. commercial launch and reentry activities, as well as the operation of non-federal launch and reentry sites within the U.S. or as carried out by U.S. citizens, as authorized by the Commercial Space Launch Act of 1984, as amended and codified at 51 U.S.C. 50901–50923. The FAA issues licenses or permits to commercial launch vehicle operators (referred to as vehicle operators or launch operators) for operation of launch and reentry vehicles. The FAA also issues licenses to commercial launch/reentry site operators for operation of commercial launch/reentry sites. An FAA license or permit is required for any commercial launch or reentry, or the operation of any commercial launch or reentry site. An FAA license or permit is not required for launch or reentry activities carried out by and for the federal government, such as NASA or Department of Defense (DoD) launches.

The FAA Office of Commercial Space Transportation issues the following types of licenses and permits, in accordance with 14 CFR parts 420, 433, 437, and 450:

- **License to Operate a Launch Site** (14 CFR part 420): A license to operate a launch site authorizes a licensee to offer its launch site to a launch operator (i.e., a person or company conducting the launch of a launch vehicle and any payload) for each launch point, launch vehicle type, and weight class identified in the license application and upon which the licensing determination is based. Issuance of a launch site operator license does not relieve a licensee of its obligation to comply with any other laws or regulations, nor does it confer any proprietary, property, or exclusive rights in the use of airspace or outer space. A launch site operator license remains in effect for five years from the date of issuance unless surrendered, suspended, or revoked before the expiration of the term and is renewable upon application by the licensee. Actual launches cannot occur from a launch site until a launch operator receives a vehicle operator license (14 CFR part 450) for the site.
- **License to Operate a Reentry Site** (14 CFR part 433): A license to operate a reentry site authorizes a licensee to offer its reentry site to a reentry vehicle operator to conduct reentry operations at the site. Issuance of a reentry site operator license does not relieve a licensee of its obligation to comply with any other laws or regulations, nor does it confer any proprietary, property, or exclusive rights in the use of airspace or outer space. Actual reentries cannot occur at a reentry site until a reentry vehicle operator receives a vehicle operator license (14 CFR part 450) for the site.
- **Experimental Permits** (14 CFR part 437): An experimental permit authorizes launch or reentry of a reusable suborbital rocket or a reusable suborbital launch vehicle for research and development, showing of compliance with requirements for obtaining a license, or crew training. The authorization includes pre- and post-flight ground operations. A suborbital rocket is a vehicle, rocket-propelled in whole or in part, intended for flight on a suborbital trajectory. A permit is an alternative to licensing, is valid for one year from the date it is issued, and the permittee may apply to renew the permit yearly.
- **Vehicle Operator License** (14 CFR part 450): A vehicle operator license authorizes a licensee to conduct one or more launches or reentries using the same vehicle or family of

vehicles. Launch includes the flight of a launch vehicle and pre- and post-flight ground operations. Reentry includes activities conducted in Earth orbit or outer space to determine reentry readiness and that are critical to ensuring public health and safety and the safety of property during reentry flight. Reentry also includes activities necessary to return the reentry vehicle, or vehicle component, to a safe condition on the ground after impact or landing. Vehicle operator licenses are valid for the period of time determined by the FAA Administrator as necessary to conduct the licensed activity, but may not exceed five years from the issuance date. The FAA may modify a vehicle operator license at any time by modifying or adding license terms and conditions to ensure compliance with the Commercial Space Launch Act and regulations.

### ***U.S. Space Force***

Under the USSF Space Systems Command, Space Launch Delta (SLD) 45 oversees launch operations at Cape Canaveral Space Force Station (CCSFS) and SLD 30 oversees launch operations at Vandenberg Space Force Base (VSFB). The USSF is the lease or license holder for the real property and ranges where launches occur from CCSFS and VSFB. The USSF uses its own launch and reentry vehicles, as well as those of commercial launch operators, to launch USSF payloads into space.

- **Space Launch Delta 45:** SLD 45 is responsible for overseeing launch operations at CCSFS (which includes multiple space launch complexes [SLC]) and ensuring range safety and supporting space missions, including launching of U.S. government and commercial payloads into orbit, from the Eastern Range (northwest Atlantic Ocean). SLD 45 also provides launch facilities and services to support NASA and commercial space operations. A USSF directive is to provide efficient means of executing national security and military policy goals. The Eastern Range operations provide the resources and activities for safe flight, range instrumentation, infrastructure, and schedule to support space and ballistic launches. The Eastern Range consists of tracking stations at CCSFS, mainland annexes, and downrange tracking stations on islands located in the Caribbean Sea and South Atlantic Ocean. SLD 45 is the primary missile and rocket launch organization for the USSF on the U.S. East Coast.
- **Space Launch Delta 30:** SLD 30, headquartered at VSFB, is responsible for overseeing launch operations at VSFB, which includes multiple SLCs. The primary mission of VSFB is to launch and track satellites destined for polar or near-polar orbit, test and evaluate America's Intercontinental Ballistic Missile systems, and support aircraft operations. SLD 30 supports West Coast launch activities for the DoD (including U.S. Air Force and Missile Defense Agency), NASA, foreign nations, and various private contractors.

### ***National Aeronautics and Space Administration***

The National Aeronautics and Space Act of 1958 established NASA. The National Aeronautics and Space Act gives NASA the responsibility for planning, directing, and conducting the nation's civilian space program, and aeronautics and aerospace research activities. It also gives NASA the authorization to enter into cooperative agreements, leases, and contracts with public and private entities in the use of NASA's services, equipment, and facilities in support of scientific research and discovery.

- **Kennedy Space Center:** Established in 1962 as the NASA Launch Operations Center, the Kennedy Space Center (KSC) has carried out launch operations for the Apollo, Skylab, Space Shuttle, and cargo and crewed launches to the International Space Station. KSC is NASA's only launch site for human spaceflight. KSC's mission is to function as a multi-user spaceport for launch operations operated by NASA and a growing number of private partners. In addition to providing all aspects of launch, landing, and recovery operations for both government and commercial launch providers, KSC also provides payload processing, testing, and integration for government and commercial partners at facilities across KSC. KSC is located adjacent to CCSFS, and the two entities work closely together to execute their missions, sharing resources, facilities, and infrastructure.
- **Wallops Flight Facility:** NASA Goddard Space Flight Center manages Wallops Flight Facility (WFF), the oldest active launch range in the continental U.S. and the only rocket testing and launch range owned and operated by NASA. For over 70 years, WFF has flown thousands of research vehicles in the quest for information on flight characteristics of launch vehicles and spacecraft, and in pursuit of knowledge of the Earth's upper atmosphere and the near space environment. WFF regularly provides launch support, range safety, and downrange tracking for the commercial space industry, either directly or through the Mid-Atlantic Regional Spaceport, which is a commercial launch site on Wallops Island licensed by the FAA and operated by the Virginia Commercial Space Flight Authority (Virginia Space). WFF also facilitates various DoD research, development, testing, and evaluation and training missions, including target and missile launches.

### Space Launch and Reentry Operations

Space launch and reentry operations cover several activities, which may vary based on the specific mission. For ease of understanding, in the following subsections we define commonly used terms in the space industry as they pertain to this consultation, overview the typical sequence of events for space operations, and break down the activities that occur at each phase.

#### *Definitions*

**Anomaly:** any condition during a licensed or permitted activity that deviates from what is standard, normal, or expected, during the verification or operation of a system, subsystem, process, facility, or support equipment.

**Breakup:** a failure resulting in a "breakup" (sometimes with an explosion) of the vehicle or stage, may be a type of anomaly or nominal (e.g., vehicle breakup during atmospheric reentry).

**Disposal:** the return or attempt to return, purposefully, a launch vehicle stage or component, not including a reentry vehicle, from Earth orbit to Earth, in a controlled manner. For example, operators may dispose of upper stages in the ocean.

**Drogue parachutes:** small, thin parachutes deployed at high altitude to gain control of the stage or vehicle component to which the parachute is attached. Part of the parachute system that facilitates recovery of spacecraft and payload fairings.

**Droneship:** also known as an autonomous spaceport drone ship, a self-propelled modified barge used to land and recover reusable boosters at sea.

**Expend:** to discard (as in, a stage, vehicle components such as fairings, or debris), often in the ocean. Usually items are expended during flight, shortly after takeoff.

**Fairing:** the nose cone of the upper stage that protects the payload during launch; it consists of two halves that separate to facilitate deployment of the payload.

**First stage:** also known as the booster, it is the bottom stage of a launch vehicle that propels the launch vehicle during launch.

**Heat shield:** a barrier that provides thermal protection for the vehicle during stage separation or during atmospheric reentry.

**High-velocity impact:** also known as a hard landing (as opposed to a soft landing, see below), where the stage impacts the ocean at terminal velocity, resulting in a breakup of the vehicle and a potential explosive event.

**Horizontal launch:** as opposed to a vertical launch, a launch (including takeoff and landing) that occurs from a runway at the launch site (similar to a commercial airplane).

**Interstage:** a structural component of the launch vehicle that connects the first stage and the second stage.

**Landing burn:** typically the final engine firing of the stage, substantially slowing down the stage for landing.

**Landing operations:** in reference to spacecraft, the reentry, splashdown, and recovery of the spacecraft.

**Launch abort test:** a test of the stage's safety system by intentionally triggering it.

**Launch operator:** person or company who conducts or will conduct the launch of a launch vehicle and any payload.

**Launch site:** location from which a launch takes place.

**Launch vehicle:** a vehicle built to operate in, or place a payload in, outer space or a suborbital rocket.

**Main parachutes:** large parachutes deployed after the drogue parachutes on a spacecraft, to provide the final slowdown of the spacecraft for splashdown.

**Nominal:** when all aspects, in reference to launch vehicle performance, systems, flight, landing, etc., occur as expected and planned.

Orbital spaceflight: where a vehicle reaches sufficient velocity to place it into orbit.

Parafoil: a large, steerable parachute, part of the system of parachutes that facilitates recovery of payload fairings.

Payload: the cargo or equipment (e.g., satellites, scientific instruments) on a vehicle or stage designed to achieve mission goals.

Pilot parachutes: small parachutes used to deploy the drogue parachutes on reentry vehicles (i.e., capsules). The pilot parachute is ejected, inflates, and then directly pulls the drogue parachute out.

Reentry: also referred to as atmospheric reentry, the return or attempt to return, purposefully, a reentry vehicle and its payload from Earth orbit or from outer space to Earth.

Reentry burn: also known as a deorbit burn, the engines fire to transition the vehicle from orbit to atmospheric reentry.

Reentry vehicle: a vehicle designed to return from Earth orbit or outer space to Earth substantially intact, and is the payload attached to the second (or upper) stage. A reusable launch vehicle that is designed to return from Earth orbit or outer space to Earth substantially intact is also a reentry vehicle. Often referred to as a spacecraft or capsule.

Reusable: as opposed to being expended, a vehicle or stage that can be recovered and used again.

Skirt ring: a structural component that may be used on launch vehicles for additional support. On SpaceX's Falcon 9 vehicle specifically, it is composed of four segments that prevent deformation of the second stage engine nozzle during stage separation.

Sonic boom: also known as overpressure, is a shock wave creating an impulsive sound that is generated during launch and landing when the vehicle reaches supersonic speed (travels faster than the speed of sound).

Space launch and reentry operations: all activities, including but not limited to pre-launch surveillance, launches, landings, expended components and debris, etc., as described in this programmatic consultation.

Splashdown: also referred to as a soft-water landing (as opposed to a hard landing or landing at terminal velocity, see above), where the vehicle or stage lands gently in the water and does not breakup.

Stage separation: During a vertical launch, where the first stage and the upper stage of the launch vehicle physically separate. During a horizontal launch, for example, when the rocket is attached to a carrier aircraft, stage separation occurs when the rocket detaches from the aircraft.

Suborbital spaceflight: where a vehicle may reach space but does not achieve sufficient velocity to place it into orbit.

Trajectory: the planned path for the vehicle's flight.

Upper stage(s): also referred to as the second stage (or third stage, if the launch vehicle has three stages), the stage on top of the first stage, which delivers a payload into orbit.

Vertical launch: as opposed to a horizontal launch, a launch (including takeoff and landing) that occurs from a launch pad, where the launch vehicle liftoff is straight up.

### ***General Sequence of Events***

Prior to launch, weather balloons, spotter aircraft, and surveillance vessels are used to ensure safe launch conditions. Pre-launch testing, such as a static fire test (firing the engines while the vehicle is bolted to the ground), may also occur. Launch vehicles may consist of multiple stages, but typically have two stages: the first stage, also referred to as the booster, and the second stage, also referred to as the upper stage or spacecraft. The second stage typically delivers the payload to its intended orbit or to outer space, and can include a reentry vehicle such as a spacecraft or capsule.

During a launch, the launch vehicle quickly gains altitude and flies along a pre-planned trajectory, often over the ocean (particularly vertical launches). At some point downrange (i.e., some point away from the launch site based on the flight trajectory), the launch vehicle reaches supersonic speeds, generating a sonic boom, and pitches over to attain its intended trajectory. Depending on the vehicle's orientation and trajectory, it is possible for the sonic boom to overlap with the marine environment. Stage separation occurs, and the first stage falls back to Earth, either being expended in the ocean or conducting controlled engine burns to decelerate the stage and land on an ocean-going barge or back at the launch site. The upper stage continues its mission and delivers the payload before reentering Earth's atmosphere and landing in the ocean. Vessels may be deployed to recover the stages or any remaining debris.

### ***Weather Balloons***

Weather balloons are deployed to measure weather data, including wind speeds, and are necessary to determine if it is safe to launch and land vehicles. Between five and 15 weather balloons are used for each launch. Weather balloons are made of latex with a radiosonde attached to each balloon. A radiosonde is an instrument that measures and transmits atmospheric data to the launch operator, and is typically the size of a half-gallon milk carton. The latex balloon typically has a diameter at launch of approximately four feet (ft; 1.2 meters [m]). When a weather balloon is deployed, it rises approximately 12–18 mi (19–29 km) into the air and then bursts. The radiosonde and shredded balloon pieces fall back to Earth and are not recovered. The radiosonde does not have a parachute and is expected to sink to the ocean floor when it lands over water.

### ***Aircraft and Vessels***

A number of spotter aircraft, including drones, and surveillance vessels are used prior to launch to ensure that designated hazard areas are clear of non-participating crafts. A combination of radar, spotter aircraft, surveillance vessels, and law enforcement vessels may be deployed prior to launch. Vessels will be up to 200 ft (61 m) in length, except barges or dronships, which may be up to 400 ft (122 m) in length. Most fixed wing aircraft operate at altitudes of 15,000 ft (4,572 m) but may drop to 1,500 ft (457 m) to obtain a call sign from a non-participating vessel.

Ports that could be used for surveillance vessel transit include all major ports on the U.S. coasts, and airports could include any airports within 50 mi (80.5 km) from the U.S. coast.

### ***Launch Sites***

Space launches are authorized to occur at several launch sites (see Table 1). The FAA, USSF, and/or NASA may receive new proposals in the future for launch operations at existing launch sites (e.g., constructing additional infrastructure or re-developing a launch site) or new launch sites that involve operations affecting the marine environment. Upon receipt of a new proposal, the lead action agency will review the proposal and coordinate with NMFS to determine if the proposed action falls within the scope of this programmatic.

**Table 1. Current launch sites**

| <b>Launch Site</b>                                                                 | <b>Location</b>  | <b>Site Operator</b>                                              |
|------------------------------------------------------------------------------------|------------------|-------------------------------------------------------------------|
| Cecil Airport                                                                      | Florida, U.S.    | Commercial – Jacksonville Aviation Authority                      |
| Cape Canaveral Space Force Station (CCSFS) – multiple space launch complexes (SLC) | Florida, U.S.    | Government – U.S. Space Force (USSF)                              |
| CCSFS – SLC-46 and SLC-20                                                          | Florida, U.S.    | Commercial – Space Florida                                        |
| Houston Spaceport (Ellington Airport)                                              | Texas, U.S.      | Commercial – Houston Airport System                               |
| Kennedy Space Center (KSC) – multiple launch complexes                             | Florida, U.S.    | Government – National Aeronautics and Space Administration (NASA) |
| KSC – Launch and Landing Facility (formerly Shuttle Landing Facility)              | Florida, U.S.    | Commercial – Space Florida                                        |
| Mojave Air and Space Port                                                          | California, U.S. | Commercial – Mojave Air and Spaceport                             |
| Pacific Spaceport Complex Alaska                                                   | Alaska, U.S.     | Commercial – Alaska Aerospace Corporation                         |
| Rocket Lab Launch Complex 1                                                        | New Zealand      | Private – Rocket Lab                                              |
| Space Coast Regional Airport                                                       | Florida, U.S.    | Commercial – Titusville-Cocoa Airport Authority                   |
| Vandenberg Space Force Base – multiple launch complexes                            | California, U.S. | Government – U.S. Space Force                                     |
| Wallops Flight Facility (WFF)                                                      | Virginia, U.S.   | Government – NASA                                                 |

| Launch Site                                              | Location       | Site Operator                                           |
|----------------------------------------------------------|----------------|---------------------------------------------------------|
| WFF – Launch Complex 0 (Mid-Atlantic Regional Spaceport) | Virginia, U.S. | Commercial – Virginia Commercial Space Flight Authority |

### ***Launch Vehicles***

Launch vehicles, commonly known as rockets, can take off either vertically from a launch pad or horizontally from a runway. There is variability in launch vehicle design. For example: the SpaceX Falcon 9 vertical launch vehicle is a two-stage launch vehicle with an interstage, which connects the first and second stages and facilitates stage separation; the United Launch Alliance's (ULA) Atlas V vertical launch vehicle consists of first stage core booster, which can be enhanced with up to five side boosters, interstage, and second stage; and the Northrop Grumman Corporation's Pegasus horizontal launch vehicle is flown to altitude by a carrier aircraft (which is considered as part of the launch vehicle) before the three-stage rocket is released and the first stage engines ignite and the rocket continues on its mission. Companies may develop upper stages, spacecraft, or capsules that launch on top of a different company's booster. NASA or DoD may conduct launches and/or reentries of vehicles that were built by the commercial space industry, and commercial launch operators may conduct operations for NASA or DoD. Launch vehicles may use a variety of fuels and propellant combinations.

Currently, a majority of launch vehicles are expendable, meaning the stages are used once and then disposed of in the ocean. Some companies (e.g., SpaceX, Blue Origin) have developed or are planning to develop reusable launch vehicles and/or stages, which can be launched, recovered, and launched again multiple times. The action agencies developed a framework of specifications for existing and future launch vehicles (based on their knowledge of the industry) to be covered under this consultation. Based on those specifications, the total engine thrust of a launch vehicle will be a maximum of 6.5 million pounds (lbs). During launch, when the launch vehicle or booster reaches supersonic speeds either during ascent or landing, a sonic boom of up to 50 pounds per square foot (psf) will be generated. This programmatic consultation covers existing and future launch vehicles whose specifications fall under those described in Table 2.

**Table 2. Launch vehicle envelope specifications**

|                                                            | Vertical launch vehicle                                                                               | Horizontal launch vehicle    |
|------------------------------------------------------------|-------------------------------------------------------------------------------------------------------|------------------------------|
| Number of stages                                           | 3                                                                                                     | 3                            |
| Vehicle length                                             | Approximately 450 ft (137.2 m)                                                                        | Approximately 80 ft (24.4 m) |
| Vehicle Diameter                                           | Approximately 23 ft (7 m)                                                                             | Approximately 10 ft (3 m)    |
| Propellant maximum amount on takeoff for solid fuel rocket | Hydroxyl-terminated Polybutadiene (HTPB): 145,000 lbs<br>High-density polyethylene (HDPE): 23,500 lbs | HTPB: 1,800 lbs              |

|                                                         | <b>Vertical launch vehicle</b>                                                                                                                                                                                                                                  | <b>Horizontal launch vehicle</b>                                                                                                     |
|---------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------|
| Propellant maximum amount on takeoff liquid fuel rocket | Liquefied natural gas (LNG): 800,000 lbs<br>LOX: 3,200,000 lbs<br>Liquid hydrogen (LH <sub>2</sub> ): 142,000 lbs<br>Rocket Propellant-1 (RP-1): 875,000 lbs<br>RP-X (potential next generation propellant; Exxsol D40, similar to mineral spirits): 45,000 lbs | Jet A fuel: 8,000 lbs<br>Nitrous oxide (N <sub>2</sub> O): 13,250 lbs<br>Hydrogen peroxide (H <sub>2</sub> O <sub>2</sub> ): 600 lbs |

ft=feet; lbs=pounds

***Launch Abort Tests***

These may include pad abort tests (triggering the abort system while the spacecraft is on the ground, in the event of an emergency at the launch pad) and launch ascent abort tests (triggering the launch abort system while the spacecraft is ascending, in the event of an emergency during launch). The latter typically involves the breakup or explosion of the first stage over the ocean and may be conducted prior to transporting crew aboard the spacecraft, to ensure safety (for example, SpaceX's Crew Dragon conducted an in-flight abort test before it carried astronauts to the International Space Station). Both types of tests may involve a waterborne landing of the spacecraft in the Atlantic Ocean. Launch abort tests typically involve a non-propulsive spacecraft landing using a system of parachutes (see more information on parachutes under the Reentry section, below). The launch operator is responsible for recovering as many pieces of debris as feasible; it is possible debris will not be recovered.

***Launch Failure Anomaly***

A launch failure anomaly is possible during any launch operation and could result in the breakup (i.e., explosion) of the booster and/or spacecraft on or near the launch pad or landing area. Anomalies may also occur during flight.

***Expendable Stages/Components/Debris***

After launch, one or more stages of the launch vehicle, or other components of the launch vehicle, may be expended (i.e., discarded) in the ocean.

During stage separation, the first stage, which has burned up most of its fuel, and upper stage physically separate. The first stage and interstage fall back to Earth and the engines of the upper stage ignite to continue on the intended mission. Some launch vehicles such as SpaceX's Falcon 9 utilize a skirt ring to prevent deformation of the upper stage engine during stage separation, which also falls back to Earth after stage separation. The skirt ring is made of four segments that are temporarily bonded to the second stage engine nozzle, so that when the second stage engines are ignited, the bond and segments are released. For a majority of launch vehicles, the first stage,

interstage, skirt ring, and heat shield will be expended in the ocean. For reusable or partially reusable (e.g., SpaceX Falcon 9 and Falcon Heavy boosters and Blue Origin boosters are reusable) launch vehicles, the first stage either lands back at the launch site or on an ocean-going barge/platform (also referred to as a dronship).

As the upper stage continues on its mission, the payload fairing, the nose cone of the upper stage that protects the payload during launch and consists of two halves, separates to facilitate deployment of the payload. Most payload fairings are expended in the ocean as they fall back to Earth. Payload fairings under this consultation may be up to 75 ft (22.9 m) in length with a diameter of 30 ft (9.1 m). Similar to reusable boosters, some companies (e.g., SpaceX) have developed the ability to recover and reuse payload fairings (for more information on fairing recovery parachutes, see Recovery section, below).

Depending on the launch vehicle and mission, other components may also be expended in the ocean. These components could include cargo and service modules (approximately 15 ft by 15 ft by 15 ft [4.6 m by 4.6 m by 4.6 m]) and interstage adapters (approximately 15 ft [4.6 m] in length and 23 ft [7 m] in diameter). Stages and components that reenter Earth's atmosphere before being expended in the ocean are expected to break up during reentry. This results in a debris field in the ocean, consisting of parts of the stages and components that were not fully destroyed during reentry.

### ***Explosive Events***

During the development of a reusable launch vehicle, the launch vehicle may experience a landing failure before being able to land back at a launch site or land on an ocean-going barge or platform. A landing failure may result in an explosive event near the surface of the ocean. Under this programmatic, explosions will only occur when related to the development of a reusable commercial launch vehicle. Landing failure explosive events, if permitted or licensed by an action agency for development of a reusable vehicle, under this programmatic may occur only under specific conditions: 1) explosive events will occur, at minimum, 11 ft (3.4 m) above the surface of the ocean; 2) the explosive weight (calculated from the amount of remaining propellant and its explosive yield) will not exceed 55,800 kilograms (123,018 lbs); and 3) explosive events will only occur in the Atlantic portion of the action area at least 100 mi (160.9 km) from any major land mass (see Figure 5).

### ***Reentry***

Reentry, splashdown, and recovery (see Recovery section, below) are the three elements of a spacecraft landing operation. During reentry, when the reentry vehicle reaches supersonic speeds, a sonic boom is generated. Reentry sonic booms for spacecraft/capsules, given the envelope of reentry vehicle specifications developed by the action agencies, may be up to 3 psf. Reentry vehicles will be a maximum of 30 ft (9.1 m) in length and 15 ft (4.6 m) in diameter. The reentry vehicle will only have residual amounts of propellant upon reentry because a majority of propellant will have been used earlier during its mission. Propellants for reentry vehicles may include hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), monomethylhydrazine (MMH), nitrogen tetroxide (NTO and N<sub>2</sub>O<sub>4</sub>), Rocket Propellant-1 (RP-1), or propene and nitrous oxide (N<sub>2</sub>O).

If landing in the ocean (versus on land), crewed (i.e., humans on board) spacecraft need to be located closer to shore for recovery and human safety. Currently, the only approved operating spacecraft for commercial crew is SpaceX's Dragon. Although FAA's reentry license allows Dragon to land in the wider North Pacific Ocean, per an agreement with FAA's Air Traffic Organization (ATO), Dragon (whether crewed or uncrewed) is only approved to land in three specific locations approximately 15.5 mi (25 km) off the coast of California (see Figure 3). A new capsule is being designed which will land at least 13.8 mi (22.2 km) from the west coast of North America, within Zone 1 or Pacific Zone 2 (see Figure 1 and Figure 2).

### ***Recovery***

As mentioned previously, SpaceX and Blue Origin have developed the ability to recover and reuse stages and fairings, and other companies are or may develop this ability in the future. First stage recovery typically consists of maneuvering the stage so that it lands back at the launch site or on a modified ocean-going barge or platform, or an autonomous spaceport dronship (or dronship, for current SpaceX operations). SpaceX dronships currently operate in the Pacific and Atlantic oceans. They are approximately 295 ft (90 m) in length, 150 ft (46 m) wide, and are able to maintain their position at sea using thruster engines. Blue Origin's dronship currently operates in the Atlantic Ocean. It is approximately 380 ft (115.8 m) in length and 150 ft (45.7 m) in width. Dronships may be up to 450 ft (137.2 m) in length and are towed to/from port and landing sites by tugs or support vessels that may be up to 200 ft (61 m) in length. During a first stage landing on a dronship, a sonic boom directly underneath and directed towards the dronship, is produced. Sonic booms generated during booster landings are typically approximately 8 psf, but may be up to 50 psf.

Fairing recovery uses a system of parachutes. Each fairing half uses one drogue parachute and one parafoil. The parachute system slows the descent of the fairing to facilitate a soft, intact splashdown of the fairing for recovery. Following reentry of the fairing into Earth's atmosphere, the drogue parachute is deployed at altitude (approximately 50,000 ft or 15,240 m) to begin the initial slowdown and to extract the parafoil. Drogue parachutes are small, thin parachutes deployed at a high altitude to gain control of the fairing at speeds that would destroy the parafoil. Each drogue parachute is made of nylon, has an area of approximately 120 square feet (ft<sup>2</sup>; 11 square meters [m<sup>2</sup>]), and has a Kevlar or Kevlar with nylon overwrap suspension line length of approximately 72 ft (22 m). Following successful deployment of the parafoil, the drogue parachute is cut away. The parafoil is made of nylon, has an area of approximately 3,000 ft<sup>2</sup> (278.7 m<sup>2</sup>), and has a nylon or nylon with Kevlar overwrap suspension line length of approximately 50 ft (15.2 m). The fairing and parafoil are then recovered by a support vessel and/or rigid-hulled inflatable boats (RHIBs; approximately 12 ft or 3.7 m in length), which are stationed in a designated safety zone near the anticipated splashdown area. The vessel then transports the fairing back to port. Recovery of the drogue parachute is attempted if weather conditions allow and the recovery team is able to obtain a visual fix on the landing location; however, that can be difficult because the drogue parachute is deployed at high altitude.

Recovery of a reentry vehicle or spacecraft is similar to fairing recovery and uses a system of parachutes. For Dragon, two drogue parachutes and four main parachutes are used during recovery. The drogue parachutes are made of nylon, each with an area of approximately 200 ft<sup>2</sup> (18.6 m<sup>2</sup>), and suspension line length of approximately 75 ft (22 m). Shortly after the drogue

parachutes are deployed, they are released and are expected to land in the ocean approximately one mile from the reentry vehicle. The main parachutes are then deployed to slow the spacecraft down to allow for a soft splashdown in the ocean or on land. The main parachutes are made of Kevlar and nylon, each with an area of approximately 10,600 ft<sup>2</sup> (984.8 m<sup>2</sup>), and suspension line length of approximately 150 ft (45.7 m). Following splashdown, an electronic locator beacon on the vehicle assists the recovery team in locating and recovering the vehicle using a pre-positioned recovery vessel of up to 200 ft (61 m) in length and RHIBs. The recovery vessel transports the vehicle back to port. Recovery of the parachutes is attempted if weather conditions allow and the recovery team is able to visually locate the parachutes. If there are crew on the reentry vehicle, they and any time-sensitive cargo will be transported to the nearest airport via helicopter. For the new capsule, a system consisting of one pilot parachute, one drogue parachute, and one main parachute will facilitate recovery. All three of these parachutes for the planned capsule are smaller than Dragon's parachutes. The pilot parachute is small, approximately 13.5 ft (4.1 m) in diameter made of nylon broad cloth, with Kevlar suspension line lengths of approximately 8.75 ft (2.7 m), and helps deploy the drogue parachute on this capsule. Like Dragon operations, recovery of the parachutes is attempted if weather conditions allow and the recovery team is able to visually locate the parachutes.

#### Action Area

*Action area* means “all areas to be affected directly or indirectly by the Federal action and not merely the immediate area involved in the action” (50 CFR §402.02). The action area is global, and is categorized by ocean basin and further subcategorized by zones (see Figures 1–7). The action area also includes U.S. waters (except for Hawai'i and the remote Pacific) from the coast out to five nautical miles (NM) or to one of the zones in Figures 2–7 (for example, in Figure 7, vessels may transit from Puerto Rico to Zone 4), within which only vessel and aircraft transit will occur.







Figure 1. Zone 1 portion of the action area (grey)




**Figure 2. Pacific portion of the action area (Zone 1 – grey; Zone 2 – dark grey; Zone 3a – white with black outline; Zone 4 – hatch; no color – outside the action area)**




**Figure 3. Subset of Pacific portion of the action area (Zone 2 – dark grey; Zone 4 – hatch; Dragon landing areas – black outline; no color – outside the action area)**




**Figure 4. Gulf portion of the action area (Zone 2 – dark grey; Zone 3 – white with black outline; no color – outside the action area)**



**Figure 5. Atlantic portion of the action area (Zone 1 – grey; Zone 2 – dark grey; Zone 3b – white with black outline; Zone 3c – transparent grey with black outline; Zone 5 – light grey; Zone 5a – hatch; Zone 5b – black; U.S. Space Force operating area – dotted; Explosive Events area – bolded black outline; no color – outside the action area)**



**Figure 6. Subset of Atlantic portion of the action area (Zone 1 – grey; Zone 2 – dark grey; Zone 3b – white with black outline; Zone 3c – transparent grey with black outline; Zone 5 – light grey; Zone 5a – hatch; Zone 5b – black; Explosive Events area – bolded black outline; no color – outside the action area)**



**Figure 7. Bahamas/Caribbean portion of the action area (Zone 1 – grey; Zone 2 – dark grey; Zone 4 – transparent grey with black outline; Zone 5 – light grey; Exuma Sound – black outline; no color – outside the action area)**

The action area also includes specific portions of the Pacific Ocean and Indian Ocean, not depicted in Figures 1–7, that correspond to the action areas of previous consultations (see Consultation History). These previous consultations are included in this programmatic because they are ongoing FAA-licensed projects. Though the activities described in those previous consultations fall under the activities in this programmatic, some portions of those previous consultations’ action areas fall outside the action area depicted in Figures 1–7. Those areas will only be used by the specific project as described in those previous consultations (see Figures 8–10).




Figure 8. Portion of the action area associated with OPR-2024-01272, Varda Space Industries, Inc. reentry operations licensed by FAA (Zone 1 – grey; Notice to Mariners – dashed line; debris areas – black outline; no color – outside the action area)





Figure 9. Portion of the action area associated with OPR-2024-02184, proposed FAA license to Sierra Space Corporation (Zone 1 – grey; Zone 2 – dark grey; Zone 3a – white; Sierra Space disposal ranges – black outline; no color – outside the action area)



**Figure 10. Portion of the action area associated with OPR-2024-01961, proposed FAA license to Boeing Company (Zone 1 – grey; Zone 2 – dark grey; Boeing debris areas – black outline; no color – outside the action area)**

### **Project Design Criteria**

Project design criteria (PDCs) are generally best management practices for a particular action. In a programmatic consultation, they are applicable to all projects implemented under the program. PDCs include measures that avoid or reduce the severity of the effects of the program on ESA-listed species and their critical habitat. The PDCs also identify planned monitoring, which is used to observe or check the progress of the actions and minimization measures over time to ensure that any measures implemented to reduce or avoid adverse effects on ESA-listed species and their critical habitats are successful.

As part of this programmatic, launch and reentry operations will be conducted by the USSF, NASA, or FAA-licensed (or permitted) commercial space operators from a launch site identified in Table 1. Operations will occur in compliance with PDCs currently implemented at these existing launch sites. Space launch operations will utilize launch vehicles that fall under the launch vehicle envelopes (Table 2).

### **General Project Design Criteria**

The following are PDCs that will be implemented in all portions of the action area:

1. All space launch and reentry operations will occur within the action area and at least 5 NM (9.3 km) from the coast of the U.S. and U.S. islands and territories (except Hawai'i). The only exception to this is vessel or aircraft transit for surveillance or recovery within 5 NM

- (9.3 km) of the coast (note this nearshore vessel and aircraft transit area is not depicted in the action area figures).
2. Stages, components, and debris will not be expended, and disposals will not occur, over coral reef areas.
  3. Space launch and reentry operations will not occur in a National Marine Sanctuary or Marine National Monument, or affect National Marine Sanctuary or Marine National Monument resources, without the appropriate authorization, permit, or consultation.
  4. In the event of a launch failure, launch operators will follow the emergency response and cleanup procedures outlined in their Hazardous Material Emergency Response Plan (or similar plan).
  5. Launch operators will instruct all personnel associated with space launch and reentry operations about ESA-listed marine species and critical habitat, and species protected under the MMPA that may be present in the operating area. Note that the Federal action agency is responsible for ensuring ESA compliance (launch operators are responsible for MMPA compliance). Launch operators will notify personnel associated with space launch and reentry operations of the civil and criminal penalties for unauthorized take of ESA-listed and MMPA-protected species.
  6. All vessel operators will be on the lookout for and attempt to avoid collision with ESA-listed and MMPA-protected species.
  7. Launch operators will provide a dedicated observer(s), defined as a biologist or person other than the vessel operator who can recognize ESA-listed and MMPA-protected species, for all in-water operations. The observer(s) is responsible for continuous monitoring of ESA-listed and MMPA-protected species with the aid of binoculars during all in-water operations (e.g., surveillance, recovery).
    - 7.1. The dedicated observer will a) alert the vessel operators to implement the appropriate vessel speed restrictions and vessel avoidance measures, and b) gather relevant data.
    - 7.2. Dedicated observers will record the date, time, GPS location, species, number of animals, distance and bearing from the vessel, direction of travel, behavior, effects to the animals from space launch and reentry operations, and other relevant information for all sightings of ESA-listed or MMPA-protected species.
    - 7.3. Dedicated observers will monitor the launch, landing, recovery areas (including areas where things are expended or disposed of, as feasible) for effects to ESA-listed and MMPA-protected species. Any effects, including injury and mortality, will be reported as noted below.
  8. When an ESA-listed or MMPA-protected species is observed, vessel operators will do the following:
    - 8.1. Maintain a minimum distance of 150 ft (45.7 m) from sea turtles.
    - 8.2. Maintain a minimum distance of 300 ft (91.4 m) from all other ESA-listed and MMPA-protected species.
    - 8.3. If the distance from the vessel to the ESA-listed or MMPA-protected species becomes less than the distance specified in these PDCs, reduce vessel speed and shift the engine to neutral. Do not engage the engines until the animals are clear of the area and transit route.
    - 8.4. Reduce vessel speed to 10 knots or less when mother/calf pairs or groups (more than two individuals) of marine mammals are observed.

- 8.5. Attempt to remain parallel to (e.g., when animals are bow-riding), or move the vessel away from, an ESA-listed or MMPA-protected species' course while the vessel is in transit and avoid excessive speed or abrupt changes in direction until the animal(s) have left the area or transit route.
9. Launch operators will immediately report any ESA-listed or MMPA-protected species collision(s), injury(ies), mortality(ies), entanglement(s), or stranding(s) to the appropriate NMFS regional contact listed below (see also <https://www.fisheries.noaa.gov/report>), and to Tanya Dobrzynski, Chief, ESA Interagency Cooperation Division, by email at [Tanya.Dobrzynski@noaa.gov](mailto:Tanya.Dobrzynski@noaa.gov), and to [nmfs.hq.esa.consultations@noaa.gov](mailto:nmfs.hq.esa.consultations@noaa.gov) with the subject line "OPR-2025-02196 – Collision, Injury, Mortality, or Stranding Report."
  - 9.1. For operations off Maine through Virginia: for marine mammals and sea turtles 866-755-6622 (also report North Atlantic right whales to the U.S. Coast Guard via VHF Channel 16 and through the [Whale Alert app](#)) and for sturgeon 978-281-9328 or [noaa.sturg911@noaa.gov](mailto:noaa.sturg911@noaa.gov)
  - 9.2. For operations off North Carolina through Texas, and the Caribbean: for marine mammals (877) WHALE-HELP (877-942-5343; also report North Atlantic right whales to the U.S. Coast Guard via VHF Channel 16 and through the [Whale Alert app](#)) and for sea turtles (844) SEA-TRTL (844-732-8785) and for sturgeon 844-788-7491 or [noaa.sturg911@noaa.gov](mailto:noaa.sturg911@noaa.gov)
  - 9.3. For operations off California, Oregon, and Washington: for marine mammals and sea turtles (877) SOS-WHALE (877-767-9425) or 866-767-6114
  - 9.4. For operations off Alaska: for marine mammals and sea turtles 877-925-7773
  - 9.5. For operations off Hawaii: for marine mammals and sea turtles 888-256-9840
  - 9.6. Off Florida, report any smalltooth sawfish sightings to (844) 4SAWFISH (844-472-9347) or via email to [sawfish@fwc.com](mailto:sawfish@fwc.com)
  - 9.7. Report any giant manta ray sightings to (727) 824-5312 or via email to [manta.ray@noaa.gov](mailto:manta.ray@noaa.gov)
10. Aircraft will maintain a baseline minimum height of 1,000 ft (304.8 m) above ESA-listed or MMPA-protected species
11. Aircraft will avoid flying in circles over any marine mammals and sea turtles, if observed, to avoid harassing behaviors.

#### ***Area-specific Project Design Criteria***

The following PDCs correspond to specific ocean basins and zones depicted in Figures 2–7 (there are no limits to the operations or number of operations in Zone 1; see Figure 1) and will be implemented in those specified areas as part of the proposed action. Dependent on mission needs, the number of annual launch and recovery operations can be variable. Table 3 below outlines the maximum annual operations expected by the action agencies that will occur in Zone 2 of the Atlantic Ocean, Gulf, and Pacific Ocean portions of the action area.

PDCs for the Pacific portion of the action area (Figure 2 and Figure 3; note no activities will occur within the Hawai'i exclusive economic zone [EEZ] or the Papahānaumokuākea National Marine Sanctuary):

- Operations in Zone 2 are limited to the number of maximum annual operations described in Table 3.

- Operations in Zone 3a are limited to vessel transit only from December 1 through April 30 to minimize impacts on ESA-listed humpback whales when listed DPS breeding aggregations are likely to be present.
- Operations in Zone 4 are limited to vessel transit and barge landings only to minimize impacts on ESA-listed species and critical habitat.
- Note that Isla Guadalupe and surrounding waters are outside the action area (i.e., no operations will occur there) to minimize impacts on the ESA-listed Guadalupe fur seal as that location is the only known breeding site for Guadalupe fur seals.
- No operations will occur over areas with sea ice cover off Alaska to minimize impacts on ESA-listed Arctic seals.
- Steller sea lion critical habitat in:
  - Alaska
    - No operations will occur within the aquatic zone that extends 20 NM (37 km) seaward in State and Federally managed waters from the baseline or basepoint of each major rookery and major haulout that is west of 144° W longitude.
    - No operations will occur within the aquatic zone that extends 3,000 ft (0.9 km) seaward in State and Federally managed waters from the baseline or basepoint of each major rookery and major haulout in Alaska east of 144° W longitude.
    - No operations will occur within the air zone that extends 3,000 ft (0.9 km) above the terrestrial zone of each major rookery and major haulout in Alaska (measured vertically from sea level).
    - No vessel operations will occur within 3 NM (5.6 km) of all Steller sea lion rookeries west of 150° W longitude.
    - In the unlikely event that debris is expected to occur over or land within the air or aquatic critical habitat zones, debris should be monitored and cleaned up if in the water. If debris lands within 3 NM (5.6 km) of a rookery west of 150° W longitude, or if debris occurs on a major rookery or major haulout (i.e., on land), notify the NMFS Alaska Regional Office and the ESA Interagency Cooperation Division at [akr.prd.section7@noaa.gov](mailto:akr.prd.section7@noaa.gov) and [nmfs.hq.esa.consultations@noaa.gov](mailto:nmfs.hq.esa.consultations@noaa.gov) with the subject line “OPR-2025-02196 – Debris in Steller sea lion critical habitat.”
  - California and Oregon
    - No operations will occur in an aquatic zone that extends 3,000 ft (0.9 km) seaward from major rookeries.
    - No operations will occur in an air zone that extends 3,000 ft (0.9 km) above the terrestrial zone of major rookeries (measured vertically from sea level).

PDCs for the Gulf portion of the action area (Figure 4):

- Operations in Zone 2 are limited to the number of maximum annual operations described in Table 3.

- Operations in Zone 3 are limited to vessel transit only to minimize impacts on ESA-listed species.
- No operations will occur within Rice's whale core distribution area.
- No operations will occur within 20 NM (37 km) of the Flower Garden Banks National Marine Sanctuary and the Florida Keys National Marine Sanctuary or listed coral reef areas.
- Vessels will slow to 10 knots or less and maintain a minimum distance of 1,500 ft (457.2 m) from Rice's whale. If a whale is observed but cannot be confirmed as a species other than a Rice's whale, the vessel operator must assume that it is a Rice's whale.
- Vessels will maintain speeds to 10 knots or less in Rice's whale proposed critical habitat.

PDCs for the Atlantic portion of the action area (Figure 5 and Figure 6):

- Operations in Zone 2 are limited to the number of maximum annual operations described in Table 3.
- Operations in Zone 3b are limited to vessel transit only, from November 1 through April 30 to minimize impacts on North Atlantic right whales during calving season.
- Operations in Zone 3c are limited to vessel transit only, from January 1 through July 31 to minimize impacts on foraging North Atlantic right whales.
- Operations in Zone 5 (off North Carolina and Florida) will not include any disposals or expended stages, expended components, or debris to minimize impacts to Northwest Atlantic Ocean DPS loggerhead sea turtle constricted migratory critical habitat and proposed North Atlantic DPS green turtle reproductive and migratory critical habitat.
- Operations in Zone 5a will not include any disposals or expended stages, expended components, or debris between November 1 through April 30 to minimize impacts to Northwest Atlantic Ocean DPS loggerhead sea turtles when they occur in high concentrations during winter months in designated critical habitat.
- Operations in Zone 5b will not include any disposals or expended stages, expended components, or debris between April 1 through November 30 to minimize impacts to Northwest Atlantic Ocean DPS loggerhead sea turtles when they occur in high densities during the breeding season in designated critical habitat.
- Vessels will slow to 10 knots or less and maintain a minimum distance of 1,500 ft (457.2 m) from North Atlantic right whales.
- Vessels will comply with the Right Whale Ship Strike Reduction Rule (50 CFR §224.105)<sup>2</sup> and reduce speeds to 10 knots or less in Seasonal Management Areas or in Right Whale Slow Zones, which are dynamic management areas established where North Atlantic right whales have been recently seen or heard (the Whale Alert app automatically notifies when entering one of these areas).
- Check various communication media for general information regarding avoiding vessel strikes and specific information regarding North Atlantic right whale sightings in the area. These include NOAA weather radio, U.S. Coast Guard NAVTEX broadcasts, and Notices to Mariners.
- Aircraft will maintain a baseline minimum height of 1,500 ft (457.2 m) above North Atlantic right whales.

<sup>2</sup> See: <http://www.fisheries.noaa.gov/pr/shipstrike/>.

- Launch abort tests will not occur in designated critical habitat for the North Atlantic right whale, Seasonal Management Areas, or Right Whale Slow Zones, and will not occur during the North Atlantic right whale winter calving season from November 1 through April 30.

Bahamas/Caribbean portion of the action area (Figure 7):

- Operations in Zone 4 are limited to vessel transit and barge landings only to minimize impacts on ESA-listed corals.
- Operations in the Exuma Sound are limited to vessel transit, barge landings, and fairing recoveries.

**Table 3. Maximum annual operations in Zone 2**

| Type of Operation                                                                      | Maximum annual operations in Atlantic Ocean Zone 2 | Maximum annual operations in Gulf Zone 2 | Maximum annual operations in Pacific Zone 2 |
|----------------------------------------------------------------------------------------|----------------------------------------------------|------------------------------------------|---------------------------------------------|
| Launches involving stages and fairings that are expended                               | 350                                                | 0                                        | 470                                         |
| Launches involving attempted recovery of stages and fairings                           | 210                                                | 0                                        | 100                                         |
| Launches involving attempted recovery of stages and expended fairings                  | 40                                                 | 0                                        | 0                                           |
| Spacecraft reentry and landing in the ocean (may involve debris expended in the ocean) | 10                                                 | 22                                       | 70                                          |
| Spacecraft reentry to a landmass with debris expended in the ocean                     | 0                                                  | 0                                        | 10                                          |
| Launch abort test                                                                      | 1                                                  | 0                                        | 0                                           |

Note: these numbers do not include operations conducting upper stage atmospheric reentry and disposing or expending the stage, components, or debris in the ocean 200 NM from land (those operations will still be included in annual reports)

### **Reporting**

The FAA, USSE, and NASA, in collaboration with launch operators, plan to prepare and submit reports to NMFS by April 1 each year and new annual reports will be produced each year activities covered under this consultation occur. Annual reports will be submitted electronically to [nmfs.hq.esa.consultations@noaa.gov](mailto:nmfs.hq.esa.consultations@noaa.gov) with the subject line “Annual Report OPR-2025-02196 Space Launch and Reentry Program.” The reports will document the action agencies’ activities

from the previous year, particularly activities that occurred in or may have affected the marine environment or ESA-listed species under NMFS's jurisdiction. See Appendix B.

Annual reports will include the following for all activities covered under this programmatic:

- 1) The dates and locations of all missions, including launch site, launch and reentry vehicles and any relevant license or permit that authorized the activities;
- 2) Contact information for the agencies and commercial entities involved in the events;
- 3) Details of launch and reentry operations that may affect the marine environment or ESA-listed species, such as stage landings at sea, activities that involve entry of materials into the marine environment, and activities that may indirectly affect the marine environment;
- 4) Dates of reentry and recovery operations if different from launch date;
- 5) Approximate locations with GPS coordinates (or last known GPS coordinates) of all landing locations, including recovered and unrecovered stages, fairings, skirt rings, parachutes, debris, and any other components that end up in the marine environment.
- 6) Information regarding vessels used during operations and transit routes, as well as aircraft activity associated with an event;
- 7) Any available information on the location and fate of unrecovered parachutes, parafoils, expended components, and debris;
- 8) Information regarding the implementation of the PDCs described above, including any relevant PDCs not implemented or issues with implementing the PDCs;
- 9) Information regarding any potential effects to ESA-listed species and critical habitat due to the activities; and
- 10) Sighting logs with observations of ESA-listed species with date, time, location, species, number of animals, distance and bearing from the vessel, direction of travel, behavior, and other relevant information.

The summary of annual aggregate activities and associated effects will allow NMFS to evaluate, among other things, whether the scope of the activities are consistent with the description of the proposed action and action area, and whether the nature and scale of the effects to ESA-listed species and critical habitat predicted continue to be valid.

#### ***Project-Specific Review***

Project-specific reviews for launch and reentry proposals that may affect an ESA-listed species or critical habitat will not require additional consultation as long as the activities are within the scope of this programmatic's identified framework, above. Action agencies may use the checklist in Appendix A to help determine whether a project falls within the scope of the programmatic framework. When projects do not fully meet those requirements, a project-specific review is required. The project lead action agency will coordinate with NMFS to determine if the scope of the project is within the scope of this programmatic, or if a separate, individual consultation is required. Examples of when an action agency should coordinate a project-specific review include: operations using a new launch site, use of a launch or reentry vehicle that is not the same as those analyzed here or does not fall under the vehicle envelopes previously described, substantial changes in technology or operations, or when activities cannot fully comply with the applicable PDCs.

If an action agency has any uncertainty that a proposed activity falls within the scope of this programmatic or needs a project-specific review, they should contact the NMFS Office of Protected Resources ESA Interagency Cooperation Division. Project-specific review requests should be sent by email to the consultation biologist and to [nmfs.hq.esa.consultations@noaa.gov](mailto:nmfs.hq.esa.consultations@noaa.gov) with the subject line “Project Specific Review Request OPR-2025-02196 Space Launch and Reentry Program” and include the following information: a project description that details the proposed operations, including details on the activities that the action agency identified do not fall under this programmatic or are unsure whether they fall under this programmatic, where and when they will occur, the number or frequency of the proposed operations, any criteria or measures that may not be fully implemented and any proposed changes to these measures or additional measures, and determination of effects to ESA-listed species and critical habitat that could result from the project. Requests for a project-specific review should be submitted at least six months in advance of the proposed activity to allow time for completion of a formal ESA section 7 consultation if one is required.

#### **SPECIES AND CRITICAL HABITAT THAT MAY BE AFFECTED BY THE PROPOSED ACTION**

Implementing regulations for the ESA require consultation on any action which may affect ESA listed species or critical habitat (50 CFR §402.14(a)). No consultation is required when an action would cause no effect. The Act envisioned all consultations to be formal, but the implementing regulations established an informal consultation process when effects are unlikely (50 CFR §402.13). Therefore, there are three general determinations when analyzing the effects of an action on listed species and critical habitat: 1) no effect, 2) may affect, likely to adversely affect (LAA; formal consultation), and 3) may affect, not likely to adversely affect (NLAA). Action agencies, prior to requesting ESA consultation, determine whether their proposed action may affect ESA-listed or proposed species or their designated or proposed critical habitat. Generally, a “no effect” determination means there is no plausible exposure or response to stressors generated by the proposed action for any ESA-listed species or critical habitat. Any scenario for which there is a plausible exposure to stressors generated by the action is considered “may affect.” For any action that “may affect” an ESA-listed species or its critical habitat, the action agency shall consult with the Services under section 7(a)(2) of the ESA.

This section identifies the ESA-listed or proposed species and designated or proposed critical habitat that occurs in the action area and thus may be affected by the stressors introduced to the action area by the proposed action. Critical habitats that occur in the action area are shown in Figures 11–27, and physical and biological features (PBFs) that may be affected by the program are discussed in the Effects Analysis. Detailed information about the biology, habitat, and conservation status of the species and critical habitat listed in Table 4 can be found in species’ status reviews, recovery plans, *Federal Register* notices, and other sources [here](#).

**Table 4. Species and critical habitat present in the action area**

| <b>Species</b>                                                 | <b>ESA Status</b>                      | <b>Critical Habitat</b>            | <b>Recovery Plan</b>    |
|----------------------------------------------------------------|----------------------------------------|------------------------------------|-------------------------|
| Beluga Whale ( <i>Delphinapterus leucas</i> ) – Cook Inlet DPS | <a href="#">E – 73 Fed. Reg. 62919</a> | <a href="#">76 Fed. Reg. 20179</a> | <a href="#">12/2016</a> |

| Species                                                                                    | ESA Status                                                    | Critical Habitat                                                         | Recovery Plan                                                   |
|--------------------------------------------------------------------------------------------|---------------------------------------------------------------|--------------------------------------------------------------------------|-----------------------------------------------------------------|
| Blue Whale ( <i>Balaenoptera musculus</i> )                                                | <a href="#">E – 35 Fed. Reg. 18319</a>                        | -- --                                                                    | <a href="#">07/1998</a><br><a href="#">11/2020</a>              |
| Bowhead Whale ( <i>Balaena mysticetus</i> )                                                | <a href="#">E – 35 Fed. Reg. 18319</a>                        | -- --                                                                    | -- --                                                           |
| Fin Whale ( <i>Balaenoptera physalus</i> )                                                 | <a href="#">E – 35 Fed. Reg. 18319</a>                        | -- --                                                                    | <a href="#">07/2010</a>                                         |
| Gray Whale ( <i>Eschrichtius robustus</i> ) – Western North Pacific DPS                    | <a href="#">E – 35 Fed. Reg. 18319</a>                        | -- --                                                                    | -- --                                                           |
| Humpback Whale ( <i>Megaptera novaeangliae</i> ) – Cape Verde Islands/Northwest Africa DPS | <a href="#">E – 81 Fed. Reg. 62259</a>                        | -- --                                                                    | <a href="#">11/1991</a>                                         |
| Humpback Whale ( <i>Megaptera novaeangliae</i> ) – Central America DPS                     | <a href="#">E – 81 Fed. Reg. 62259</a>                        | <a href="#">86 Fed. Reg. 21082</a>                                       | <a href="#">11/1991</a><br><a href="#">06/2022</a><br>(Outline) |
| Humpback Whale ( <i>Megaptera novaeangliae</i> ) – Mexico DPS                              | <a href="#">T – 81 Fed. Reg. 62259</a>                        | <a href="#">86 Fed. Reg. 21082</a>                                       | <a href="#">11/1991</a><br><a href="#">06/2022</a><br>(Outline) |
| Humpback Whale ( <i>Megaptera novaeangliae</i> ) – Western North Pacific DPS               | <a href="#">E – 81 Fed. Reg. 62259</a>                        | <a href="#">86 Fed. Reg. 21082</a>                                       | <a href="#">11/1991</a><br><a href="#">06/2022</a><br>(Outline) |
| Killer Whale ( <i>Orcinus orca</i> ) – Southern Resident DPS                               | <a href="#">E – 70 Fed. Reg. 69903 and 80 Fed. Reg. 7380</a>  | <a href="#">71 Fed. Reg. 69054</a><br><a href="#">86 Fed. Reg. 41668</a> | <a href="#">01/2008</a>                                         |
| North Atlantic Right Whale ( <i>Eubalaena glacialis</i> )                                  | <a href="#">E – 73 Fed. Reg. 12024</a>                        | <a href="#">81 Fed. Reg. 4837</a>                                        | <a href="#">08/2004</a>                                         |
| North Pacific Right Whale ( <i>Eubalaena japonica</i> )                                    | <a href="#">E – 73 Fed. Reg. 12024</a>                        | <a href="#">73 Fed. Reg. 19000</a>                                       | <a href="#">06/2013</a>                                         |
| Rice's Whale ( <i>Balaenoptera ricei</i> )                                                 | <a href="#">E – 84 Fed. Reg. 15446 and 86 Fed. Reg. 47022</a> | <a href="#">88 Fed. Reg. 47453</a> (Proposed)                            | <a href="#">09/2020</a><br>(Outline)                            |
| Sei Whale ( <i>Balaenoptera borealis</i> )                                                 | <a href="#">E – 35 Fed. Reg. 18319</a>                        | -- --                                                                    | <a href="#">12/2011</a>                                         |
| Southern Right Whale ( <i>Eubalaena australis</i> )                                        | <a href="#">E – 35 Fed. Reg. 18319</a>                        | -- --                                                                    |                                                                 |
| Sperm Whale ( <i>Physeter macrocephalus</i> )                                              | <a href="#">E – 35 Fed. Reg. 18319</a>                        | -- --                                                                    | <a href="#">12/2010</a>                                         |
| Bearded Seal ( <i>Erignathus barbatus nauticus</i> ) – Beringia DPS                        | <a href="#">T – 77 Fed. Reg. 76739</a>                        | <a href="#">87 Fed. Reg. 19180</a>                                       | -- --                                                           |
| Guadalupe Fur Seal ( <i>Arctocephalus townsendi</i> )                                      | <a href="#">T – 50 Fed. Reg. 51252</a>                        | -- --                                                                    | -- --                                                           |
| Ringed Seal ( <i>Phoca hispida hispida</i> ) – Arctic subspecies                           | <a href="#">T – 77 Fed. Reg. 76706</a>                        | <a href="#">87 Fed. Reg. 19232</a>                                       | -- --                                                           |

| Species                                                                   | ESA Status                             | Critical Habitat                                                                    | Recovery Plan                                                                                                    |
|---------------------------------------------------------------------------|----------------------------------------|-------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|
| Steller Sea Lion ( <i>Eumetopias jubatus</i> ) – Western DPS              | <a href="#">E – 55 Fed. Reg. 49204</a> | <a href="#">58 Fed. Reg. 45269</a>                                                  | <a href="#">03/2008</a>                                                                                          |
| Green Sea Turtle ( <i>Chelonia mydas</i> ) – Central North Pacific DPS    | <a href="#">T – 81 Fed. Reg. 20057</a> | <a href="#">88 Fed. Reg. 46572</a> (Proposed)*                                      | <a href="#">01/1998</a>                                                                                          |
| Green Sea Turtle ( <i>Chelonia mydas</i> ) – Central South Pacific DPS    | <a href="#">E – 81 Fed. Reg. 20057</a> | <a href="#">88 Fed. Reg. 46572</a> (Proposed)*                                      | <a href="#">01/1998</a>                                                                                          |
| Green Sea Turtle ( <i>Chelonia mydas</i> ) – Central West Pacific DPS     | <a href="#">E – 81 Fed. Reg. 20057</a> | <a href="#">88 Fed. Reg. 46572</a> (Proposed)*                                      | <a href="#">01/1998</a>                                                                                          |
| Green Sea Turtle ( <i>Chelonia mydas</i> ) – East Indian-West Pacific DPS | <a href="#">T – 81 Fed. Reg. 20057</a> | -- --                                                                               | -- --                                                                                                            |
| Green Sea Turtle ( <i>Chelonia mydas</i> ) – East Pacific DPS             | <a href="#">T – 81 Fed. Reg. 20057</a> | <a href="#">88 Fed. Reg. 46572</a> (Proposed)                                       | <a href="#">01/1998</a>                                                                                          |
| Green Sea Turtle ( <i>Chelonia mydas</i> ) – North Atlantic DPS           | <a href="#">T – 81 Fed. Reg. 20057</a> | <a href="#">63 Fed. Reg. 46693</a><br><a href="#">88 Fed. Reg. 46572</a> (Proposed) | <a href="#">10/1991</a> – U.S. Atlantic                                                                          |
| Green Sea Turtle ( <i>Chelonia mydas</i> ) – North Indian DPS             | <a href="#">T – 81 Fed. Reg. 20057</a> | -- --                                                                               | -- --                                                                                                            |
| Green Sea Turtle ( <i>Chelonia mydas</i> ) – South Atlantic DPS           | <a href="#">T – 81 Fed. Reg. 20057</a> | <a href="#">88 Fed. Reg. 46572</a> (Proposed)                                       | -- --                                                                                                            |
| Green Sea Turtle ( <i>Chelonia mydas</i> ) – Southwest Indian DPS         | <a href="#">T – 81 Fed. Reg. 20057</a> | -- --                                                                               | -- --                                                                                                            |
| Green Sea Turtle ( <i>Chelonia mydas</i> ) – Southwest Pacific DPS        | <a href="#">T – 81 Fed. Reg. 20057</a> | -- --                                                                               | -- --                                                                                                            |
| Hawksbill Sea Turtle ( <i>Eretmochelys imbricata</i> )                    | <a href="#">E – 35 Fed. Reg. 8491</a>  | <a href="#">63 Fed. Reg. 46693</a>                                                  | <a href="#">08/1992</a> – U.S. Caribbean, Atlantic, and Gulf of Mexico<br><a href="#">05/1998</a> – U.S. Pacific |
| Kemp's Ridley Sea Turtle ( <i>Lepidochelys kempii</i> )                   | <a href="#">E – 35 Fed. Reg. 18319</a> | -- --                                                                               | <a href="#">03/2010</a> – U.S. Caribbean, Atlantic, and Gulf of Mexico<br><a href="#">09/2011</a>                |
| Leatherback Sea Turtle ( <i>Dermochelys coriacea</i> )                    | <a href="#">E – 35 Fed. Reg. 8491</a>  | <a href="#">44 Fed. Reg. 17710</a><br><a href="#">77 Fed. Reg. 4170</a>             | <a href="#">10/1991</a> – U.S. Caribbean, Atlantic, and Gulf of Mexico                                           |

| Species                                                                                             | ESA Status                                                    | Critical Habitat                   | Recovery Plan                          |
|-----------------------------------------------------------------------------------------------------|---------------------------------------------------------------|------------------------------------|----------------------------------------|
|                                                                                                     |                                                               |                                    | <a href="#">05/1998</a> – U.S. Pacific |
| Loggerhead Sea Turtle ( <i>Caretta caretta</i> ) – North Indian Ocean DPS                           | <a href="#">E – 76 Fed. Reg. 58868</a>                        | -- --                              | -- --                                  |
| Loggerhead Sea Turtle ( <i>Caretta caretta</i> ) – North Pacific Ocean DPS                          | <a href="#">E – 76 Fed. Reg. 58868</a>                        | -- --                              | <a href="#">01/1998</a> – U.S. Pacific |
| Loggerhead Sea Turtle ( <i>Caretta caretta</i> ) – Northeast Atlantic Ocean DPS                     | <a href="#">E – 76 Fed. Reg. 58868</a>                        | -- --                              | -- --                                  |
| Loggerhead Sea Turtle ( <i>Caretta caretta</i> ) – Northwest Atlantic Ocean DPS                     | <a href="#">T – 76 Fed. Reg. 58868</a>                        | <a href="#">79 Fed. Reg. 39855</a> | <a href="#">12/2008</a>                |
| Loggerhead Sea Turtle ( <i>Caretta caretta</i> ) – South Atlantic Ocean DPS                         | <a href="#">T – 76 Fed. Reg. 58868</a>                        | -- --                              | -- --                                  |
| Loggerhead Sea Turtle ( <i>Caretta caretta</i> ) – South Pacific Ocean DPS                          | <a href="#">E – 76 Fed. Reg. 58868</a>                        | -- --                              | -- --                                  |
| Loggerhead Sea Turtle ( <i>Caretta caretta</i> ) – Southeast Indo-Pacific Ocean DPS                 | <a href="#">T – 76 Fed. Reg. 58868</a>                        | -- --                              | -- --                                  |
| Loggerhead Sea Turtle ( <i>Caretta caretta</i> ) – Southwest Indian Ocean DPS                       | <a href="#">T – 76 Fed. Reg. 58868</a>                        | -- --                              | -- --                                  |
| Olive Ridley Sea Turtle ( <i>Lepidochelys olivacea</i> ) – All Other Areas                          | <a href="#">T – 43 Fed. Reg. 32800</a>                        | -- --                              | -- --                                  |
| Olive Ridley Sea Turtle ( <i>Lepidochelys olivacea</i> ) – Mexico's Pacific Coast Breeding Colonies | <a href="#">E – 43 Fed. Reg. 32800</a>                        | -- --                              | <a href="#">63 Fed. Reg. 28359</a>     |
| Atlantic Salmon ( <i>Salmo salar</i> ) – Gulf of Maine DPS                                          | <a href="#">E – 65 Fed. Reg. 69459 and 74 Fed. Reg. 29344</a> | <a href="#">74 Fed. Reg. 39903</a> | <a href="#">02/2019</a>                |
| Atlantic Sturgeon ( <i>Acipenser oxyrinchus oxyrinchus</i> ) – Carolina DPS                         | <a href="#">E – 77 Fed. Reg. 5913</a>                         | <a href="#">82 Fed. Reg. 39160</a> | <a href="#">02/2012</a> (Outline)      |
| Atlantic Sturgeon ( <i>Acipenser oxyrinchus oxyrinchus</i> ) – Chesapeake Bay DPS                   | <a href="#">E – 77 Fed. Reg. 5880</a>                         | <a href="#">82 Fed. Reg. 39160</a> | <a href="#">02/2012</a> (Outline)      |
| Atlantic Sturgeon ( <i>Acipenser oxyrinchus oxyrinchus</i> ) – Gulf of Maine DPS                    | <a href="#">T – 77 Fed. Reg. 5879</a>                         | <a href="#">82 Fed. Reg. 39160</a> | <a href="#">02/2012</a> (Outline)      |

| <b>Species</b>                                                                           | <b>ESA Status</b>                                            | <b>Critical Habitat</b>            | <b>Recovery Plan</b>                 |
|------------------------------------------------------------------------------------------|--------------------------------------------------------------|------------------------------------|--------------------------------------|
| Atlantic Sturgeon ( <i>Acipenser oxyrinchus oxyrinchus</i> ) – New York Bight DPS        | <a href="#">E – 77 Fed. Reg. 5879</a>                        | <a href="#">82 Fed. Reg. 39160</a> | <a href="#">02/2012</a><br>(Outline) |
| Atlantic Sturgeon ( <i>Acipenser oxyrinchus oxyrinchus</i> ) – South Atlantic DPS        | <a href="#">E – 77 Fed. Reg. 5913</a>                        | <a href="#">82 Fed. Reg. 39160</a> | <a href="#">02/2012</a><br>(Outline) |
| Bocaccio ( <i>Sebastes paucispinis</i> ) – Puget Sound/Georgia Basin DPS                 | <a href="#">E – 75 Fed. Reg. 22276 and 82 Fed. Reg. 7711</a> | <a href="#">79 Fed. Reg. 68042</a> | <a href="#">10/2017</a>              |
| Chinook Salmon ( <i>Oncorhynchus tshawytscha</i> ) – California Coastal ESU              | <a href="#">T – 70 Fed. Reg. 37160</a>                       | <a href="#">70 Fed. Reg. 52488</a> | <a href="#">10/2016</a>              |
| Chinook Salmon ( <i>Oncorhynchus tshawytscha</i> ) – Central Valley Spring-Run ESU       | <a href="#">T – 70 Fed. Reg. 37160</a>                       | <a href="#">70 Fed. Reg. 52488</a> | <a href="#">07/2014</a>              |
| Chinook Salmon ( <i>Oncorhynchus tshawytscha</i> ) – Lower Columbia River ESU            | <a href="#">T – 70 Fed. Reg. 37160</a>                       | <a href="#">70 Fed. Reg. 52629</a> | <a href="#">06/2013</a>              |
| Chinook Salmon ( <i>Oncorhynchus tshawytscha</i> ) – Puget Sound ESU                     | <a href="#">T – 70 Fed. Reg. 37160</a>                       | <a href="#">70 Fed. Reg. 52629</a> | <a href="#">01/2007</a>              |
| Chinook Salmon ( <i>Oncorhynchus tshawytscha</i> ) – Sacramento River Winter-Run ESU     | <a href="#">E – 70 Fed. Reg. 37160</a>                       | <a href="#">58 Fed. Reg. 33212</a> | <a href="#">07/2014</a>              |
| Chinook Salmon ( <i>Oncorhynchus tshawytscha</i> ) – Snake River Fall-Run ESU            | <a href="#">T – 70 Fed. Reg. 37160</a>                       | <a href="#">58 Fed. Reg. 68543</a> | <a href="#">11/2017</a>              |
| Chinook Salmon ( <i>Oncorhynchus tshawytscha</i> ) – Snake River Spring/Summer-Run ESU   | <a href="#">T – 70 Fed. Reg. 37160</a>                       | <a href="#">64 Fed. Reg. 57399</a> | <a href="#">11/2017</a>              |
| Chinook Salmon ( <i>Oncorhynchus tshawytscha</i> ) – Upper Columbia River Spring-Run ESU | <a href="#">E – 70 Fed. Reg. 37160</a>                       | <a href="#">70 Fed. Reg. 52629</a> | <a href="#">08/2007</a>              |
| Chinook Salmon ( <i>Oncorhynchus tshawytscha</i> ) – Upper Willamette River ESU          | <a href="#">T – 70 Fed. Reg. 37160</a>                       | <a href="#">70 Fed. Reg. 52629</a> | <a href="#">08/2011</a>              |
| Chum Salmon ( <i>Oncorhynchus keta</i> ) – Columbia River ESU                            | <a href="#">T – 70 Fed. Reg. 37160</a>                       | <a href="#">70 Fed. Reg. 52629</a> | <a href="#">06/2013</a>              |
| Chum Salmon ( <i>Oncorhynchus keta</i> ) – Hood Canal Summer-Run ESU                     | <a href="#">T – 70 Fed. Reg. 37160</a>                       | <a href="#">70 Fed. Reg. 52629</a> | <a href="#">11/2005</a>              |

| <b>Species</b>                                                                                   | <b>ESA Status</b>                      | <b>Critical Habitat</b>            | <b>Recovery Plan</b>              |
|--------------------------------------------------------------------------------------------------|----------------------------------------|------------------------------------|-----------------------------------|
| Coho Salmon ( <i>Oncorhynchus kisutch</i> ) – Central California Coast ESU                       | <a href="#">E – 70 Fed. Reg. 37160</a> | <a href="#">64 Fed. Reg. 24049</a> | <a href="#">09/2012</a>           |
| Coho Salmon ( <i>Oncorhynchus kisutch</i> ) – Lower Columbia River ESU                           | <a href="#">T – 70 Fed. Reg. 37160</a> | <a href="#">81 Fed. Reg. 9251</a>  | <a href="#">06/2013</a>           |
| Coho Salmon ( <i>Oncorhynchus kisutch</i> ) – Oregon Coast ESU                                   | <a href="#">T – 73 Fed. Reg. 7816</a>  | <a href="#">73 Fed. Reg. 7816</a>  | <a href="#">12/2016</a>           |
| Coho Salmon ( <i>Oncorhynchus kisutch</i> ) – Southern Oregon and Northern California Coasts ESU | <a href="#">T – 70 Fed. Reg. 37160</a> | <a href="#">64 Fed. Reg. 24049</a> | <a href="#">09/2014</a>           |
| Eulachon ( <i>Thaleichthys pacificus</i> ) – Southern DPS                                        | <a href="#">T – 75 Fed. Reg. 13012</a> | <a href="#">76 Fed. Reg. 65323</a> | <a href="#">09/2017</a>           |
| Giant Manta Ray ( <i>Manta birostris</i> )                                                       | <a href="#">T – 83 Fed. Reg. 2916</a>  | -- --                              | <a href="#">12/2019 (Outline)</a> |
| Green Sturgeon ( <i>Acipenser medirostris</i> ) – Southern DPS                                   | <a href="#">T – 71 Fed. Reg. 17757</a> | <a href="#">74 Fed. Reg. 52300</a> | <a href="#">08/2018</a>           |
| Gulf Grouper ( <i>Mycteroperca jordani</i> )                                                     | <a href="#">E – 81 Fed. Reg. 72545</a> | -- --                              | -- --                             |
| Gulf Sturgeon ( <i>Acipenser oxyrinchus desotoi</i> )                                            | <a href="#">T – 56 Fed. Reg. 49653</a> | <a href="#">68 Fed. Reg. 13370</a> | <a href="#">09/1995</a>           |
| Nassau Grouper ( <i>Epinephelus striatus</i> )                                                   | <a href="#">T – 81 Fed. Reg. 42268</a> | <a href="#">89 Fed. Reg. 126</a>   | <a href="#">08/2018 (Outline)</a> |
| Oceanic Whitetip Shark ( <i>Carcharhinus longimanus</i> )                                        | <a href="#">T – 83 Fed. Reg. 4153</a>  | -- --                              | <a href="#">07/2024</a>           |
| Scalloped Hammerhead Shark ( <i>Sphyrna lewini</i> ) – Central and Southwest Atlantic DPS        | <a href="#">T – 79 Fed. Reg. 38213</a> | -- --                              | -- --                             |
| Scalloped Hammerhead Shark ( <i>Sphyrna lewini</i> ) – Eastern Atlantic DPS                      | <a href="#">E – 79 Fed. Reg. 38213</a> | -- --                              | -- --                             |
| Scalloped Hammerhead Shark ( <i>Sphyrna lewini</i> ) – Eastern Pacific DPS                       | <a href="#">E – 79 Fed. Reg. 38213</a> | -- --                              | -- --                             |
| Shortnose Sturgeon ( <i>Acipenser brevirostrum</i> )                                             | <a href="#">E – 32 Fed. Reg. 4001</a>  | -- --                              | <a href="#">12/1998</a>           |
| Smalltooth Sawfish ( <i>Pristis pectinata</i> ) – U.S. portion of range DPS                      | <a href="#">E – 68 Fed. Reg. 15674</a> | <a href="#">74 Fed. Reg. 45353</a> | <a href="#">01/2009</a>           |
| Smalltooth Sawfish ( <i>Pristis pectinata</i> ) – Non-U.S. portion of range DPS                  | <a href="#">E - 79 Fed. Reg. 73977</a> | -- --                              | -- --                             |

| <b>Species</b>                                                                      | <b>ESA Status</b>                                            | <b>Critical Habitat</b>            | <b>Recovery Plan</b>                 |
|-------------------------------------------------------------------------------------|--------------------------------------------------------------|------------------------------------|--------------------------------------|
| Sockeye Salmon ( <i>Oncorhynchus nerka</i> ) – Ozette Lake ESU                      | <a href="#">T – 70 Fed. Reg. 37160</a>                       | <a href="#">70 Fed. Reg. 52630</a> | <a href="#">05/2009</a>              |
| Sockeye Salmon ( <i>Oncorhynchus nerka</i> ) – Snake River ESU                      | <a href="#">E – 70 Fed. Reg. 37160</a>                       | <a href="#">58 Fed. Reg. 68543</a> | <a href="#">06/2015</a>              |
| Steelhead Trout ( <i>Oncorhynchus mykiss</i> ) – California Central Valley DPS      | <a href="#">T – 71 Fed. Reg. 834</a>                         | <a href="#">70 Fed. Reg. 52487</a> | <a href="#">07/2014</a>              |
| Steelhead Trout ( <i>Oncorhynchus mykiss</i> ) – Central California Coast DPS       | <a href="#">T – 71 Fed. Reg. 834</a>                         | <a href="#">70 Fed. Reg. 52487</a> | <a href="#">10/2016</a>              |
| Steelhead Trout ( <i>Oncorhynchus mykiss</i> ) – Lower Columbia River DPS           | <a href="#">T – 71 Fed. Reg. 834</a>                         | <a href="#">70 Fed. Reg. 52629</a> | <a href="#">06/2013</a>              |
| Steelhead Trout ( <i>Oncorhynchus mykiss</i> ) – Middle Columbia River DPS          | <a href="#">T – 71 Fed. Reg. 834</a>                         | <a href="#">70 Fed. Reg. 52629</a> | <a href="#">11/2009</a>              |
| Steelhead Trout ( <i>Oncorhynchus mykiss</i> ) – Northern California DPS            | <a href="#">T – 71 Fed. Reg. 834</a>                         | <a href="#">70 Fed. Reg. 52487</a> | <a href="#">10/2016</a>              |
| Steelhead Trout ( <i>Oncorhynchus mykiss</i> ) – Puget Sound DPS                    | <a href="#">T – 72 Fed. Reg. 26722</a>                       | <a href="#">81 Fed. Reg. 9251</a>  | <a href="#">12/2019</a>              |
| Steelhead Trout ( <i>Oncorhynchus mykiss</i> ) – Snake River Basin DPS              | <a href="#">T – 71 Fed. Reg. 834</a>                         | <a href="#">70 Fed. Reg. 52629</a> | <a href="#">11/2017</a>              |
| Steelhead Trout ( <i>Oncorhynchus mykiss</i> ) – South-Central California Coast DPS | <a href="#">T – 71 Fed. Reg. 834</a>                         | <a href="#">70 Fed. Reg. 52487</a> | <a href="#">12/2013</a>              |
| Steelhead Trout ( <i>Oncorhynchus mykiss</i> ) – Southern California DPS            | <a href="#">E – 71 Fed. Reg. 834</a>                         | <a href="#">70 Fed. Reg. 52487</a> | <a href="#">01/2012</a>              |
| Steelhead Trout ( <i>Oncorhynchus mykiss</i> ) – Upper Columbia River DPS           | <a href="#">T – 71 Fed. Reg. 834</a>                         | <a href="#">70 Fed. Reg. 52629</a> | <a href="#">08/2007</a>              |
| Steelhead Trout ( <i>Oncorhynchus mykiss</i> ) – Upper Willamette River DPS         | <a href="#">T – 71 Fed. Reg. 834</a>                         | <a href="#">70 Fed. Reg. 52629</a> | <a href="#">08/2011</a>              |
| Yelloweye Rockfish ( <i>Sebastes ruberrimus</i> ) – Puget Sound/Georgia Basin DPS   | <a href="#">T – 75 Fed. Reg. 22276 and 82 Fed. Reg. 7711</a> | <a href="#">79 Fed. Reg. 68041</a> | <a href="#">10/2017</a>              |
| Black Abalone ( <i>Haliotis cracherodii</i> )                                       | <a href="#">E – 74 Fed. Reg. 1937</a>                        | <a href="#">76 Fed. Reg. 66805</a> | <a href="#">11/2020</a>              |
| Boulder Star Coral ( <i>Orbicella franksi</i> )                                     | <a href="#">T – 79 Fed. Reg. 53851</a>                       | <a href="#">88 Fed. Reg. 54026</a> | <a href="#">03/2015</a><br>(Outline) |

| Species                                                | ESA Status                                        | Critical Habitat                   | Recovery Plan                        |
|--------------------------------------------------------|---------------------------------------------------|------------------------------------|--------------------------------------|
| Chambered Nautilus ( <i>Nautilus pompilius</i> )       | <a href="#">T – 83 Fed. Reg. 48976</a>            | -- --                              | -- --                                |
| Elkhorn Coral ( <i>Acropora palmata</i> )              | <a href="#">T – 79 Fed. Reg. 53851</a>            | <a href="#">73 Fed. Reg. 72210</a> | <a href="#">03/2015</a>              |
| Lobed Star Coral ( <i>Orbicella annularis</i> )        | <a href="#">T – 79 Fed. Reg. 53851</a>            | <a href="#">88 Fed. Reg. 54026</a> | <a href="#">03/2015</a><br>(Outline) |
| Mountainous Star Coral ( <i>Orbicella faveolata</i> )  | <a href="#">T – 79 Fed. Reg. 53851</a>            | <a href="#">88 Fed. Reg. 54026</a> | <a href="#">03/2015</a><br>(Outline) |
| Pillar Coral ( <i>Dendrogyra cylindrus</i> )           | <a href="#">E – 89 Fed. Reg. 101993</a>           | <a href="#">88 Fed. Reg. 54026</a> | <a href="#">03/2015</a><br>(Outline) |
| Queen conch ( <i>Aliger gigas</i> )                    | <a href="#">T – 89 Fed. Reg. 11208</a>            | -- --                              | -- --                                |
| Rough Cactus Coral ( <i>Mycetophyllia ferox</i> )      | <a href="#">T – 79 Fed. Reg. 53851</a>            | <a href="#">88 Fed. Reg. 54026</a> | <a href="#">03/2015</a><br>(Outline) |
| Staghorn Coral ( <i>Acropora cervicornis</i> )         | <a href="#">T – 79 Fed. Reg. 53851</a>            | <a href="#">73 Fed. Reg. 72210</a> | <a href="#">03/2015</a>              |
| Sunflower Sea Star ( <i>Pycnopodia helianthoides</i> ) | <a href="#">T – 88 Fed. Reg. 16212</a> (Proposed) | -- --                              | -- --                                |
| White Abalone ( <i>Haliotis sorenseni</i> )            | <a href="#">E – 66 Fed. Reg. 29046</a>            | -- --                              | <a href="#">10/2008</a>              |

DPS=distinct population segment; ESU=evolutionarily significant unit; E=endangered; T=threatened; Fed. Reg.=Federal Register

\* = not in action area

#### ESA-Listed Marine Mammals in the Action Area

Blue whales, fin whales, and sei whales are widely distributed across the globe in all major oceans. All of these species typically winter at low latitudes, where they mate, calve, and nurse, and summer at high latitudes, where they feed. They are most common in offshore continental shelf and slope waters that support productive zooplankton blooms.

Humpback whales are also widely distributed and winter at low latitudes, where they calve and nurse, and summer at high latitudes, where they feed. The Western North Pacific DPS of humpback whales breeds/winters off Okinawa and the Philippines, and they migrate to northern feeding grounds primarily off the Russian coast. Those areas are outside of the action area of this programmatic; however, the Western North Pacific DPS of humpback whales also feeds near the Aleutian Islands and the Gulf of Alaska, which is within the action area (81 Fed. Reg. 62259). The Mexico DPS of humpback whales breeds along the Pacific coast of mainland Mexico, but also near the Revillagigedo Archipelago, which overlaps the action area, and feeds in the action area across a broad geographic range from California to the Aleutian Islands (81 Fed. Reg. 62259). The Central America DPS of humpback whales breeds along the Pacific coast of Central America and feeds in the action area almost exclusively offshore of California and Oregon (81 Fed. Reg. 62259). The Cape Verde Islands/Northwest Africa DPS consists of humpback whales primarily known to breed and calve around the Cape Verde Islands, with feeding grounds near Iceland and Norway (81 Fed. Reg. 62259).

Bowhead whales reside almost exclusively in Arctic and subarctic waters experiencing seasonal sea ice coverage, primarily between 60° and 75° north latitude. They spend the winter associated with the southern limit of the pack ice and move north as the sea ice breaks up and recedes during spring. The Western Arctic stock is found in waters around Alaska; Bering, Chukchi, and Beaufort seas (Rugh and Shelden 2009).

The Western North Pacific gray whales tend to feed near the bottom in productive waters closer to shore. Some Western North Pacific gray whales winter in the action area on the west coast of North America, while most others migrate south to winter in waters off Japan and China and summer in the Okhotsk Sea off northeast Sakhalin Island, Russia, and off southeastern Kamchatka in the Bering Sea (Burdin et al. 2013).

North Pacific right whales mostly inhabit coastal and continental shelf waters in the North Pacific Ocean. They have been observed in temperate latitudes during winter off Japan (outside the action area), California, and Mexico where they likely calve and nurse (73 Fed. Reg. 12024). In the summer, they feed on large concentrations of zooplankton in sub-polar waters around Alaska.

Southern right whales are found in the Southern Hemisphere worldwide from temperate to polar waters, favoring shallow waters less than 65.6 ft (20 m) deep. Southern right whales migrate between winter breeding areas in coastal waters of the South Atlantic, Pacific, and Indian Oceans from May to December and offshore summer (January to April) foraging locations in the Subtropical and Antarctic Convergence zones (NMFS 2015).

The North Atlantic right whale is primarily found in the western North Atlantic Ocean from shallow coastal water breeding grounds in temperate latitudes off the coast of the southeastern U.S. during the winter, to feeding in summer off the northeastern U.S. and up to Canadian waters (Waring and et al. 2016).

Rice's whales (formerly the Gulf of Mexico Bryde's whale) live year-round in the Gulf of America (formerly Gulf of Mexico) and are the only resident baleen whale in the Gulf. Most sightings of Rice's whales have been in waters 328.1–1,312.3 ft (100–400 m) deep along the continental shelf break, more often in a biologically important area of the northeastern Gulf near De Soto Canyon (Rosel et al. 2021). Sightings have also been documented, albeit less frequently, in the western Gulf, offshore of Louisiana and Texas.

The sperm whale is widely distributed globally, found in all major oceans. Sperm whales mostly inhabit areas with a water depth of 1,968 ft (600 m) or more and are uncommon in waters less than 984 ft (300 m) deep. They winter at low latitudes, where they calve and nurse, and summer at high latitudes, where they feed primarily on squid and demersal fish.

The Southern Resident DPS killer whale is found along the Pacific Coast of the U.S. and Canada. Southern Resident killer whales occur in the inland waterways of Puget Sound, the Strait of Juan de Fuca, and the Southern Georgia Strait during the spring, summer, and fall. During the winter, they move out into coastal waters primarily off Oregon, Washington, California, and British Columbia.

Cook Inlet DPS beluga whales reside in Cook Inlet year-round, which makes them geographically and genetically isolated from other beluga whale stocks (Allen et al. 2011). Within Cook Inlet, they generally occur in shallow, coastal waters, often in water barely deep enough to cover their bodies (Harrison and Ridgway 1981).

Bearded seals are boreoarctic with a circumpolar distribution and are closely associated with sea ice. Bearded seals inhabit circumpolar Arctic and sub-Arctic waters that are relatively shallow (typically less than about 650 ft or 198 m deep) and seasonally ice-covered. Beringia DPS bearded seals are found in the continental shelf waters throughout the Eastern Siberian, Chukchi, Beaufort, and Bering Seas. Bearded seals generally move north in late spring and summer as the ice melts and retreats, and then south in the fall as sea ice freezes and expands; however, some remain near the coasts during the summer and early fall.


Ringed seals have a circumpolar distribution. The Arctic subspecies of ringed seals are widely distributed throughout the Arctic Ocean, in waters of Russia, Canada, Greenland, Finland, and the U.S. In U.S. waters, Arctic ringed seals are found around Alaska in the Bering, Chukchi, and Beaufort Seas. Most seals move seasonally, following the extent of the sea ice.

The Western DPS Steller sea lions reside in the central and western Gulf of Alaska, the Aleutian Islands, as well as coastal portions of Japan and Russia (the latter of which are not in the action area). Western DPS Steller sea lions typically forage in coastal waters on the continental shelf, but they sometimes forage in deeper continental slope and pelagic waters, especially in the non-breeding season.

Guadalupe fur seals breed mainly on Guadalupe Island with another smaller breeding colony in the San Benito Archipelago, Baja California, Mexico (Belcher and T.E. Lee 2002). Guadalupe fur seals feed mainly on squid species (Esperon-Rodriguez and Gallo-Reynoso 2013) with foraging trips that can last between four to 24 days (average of 14 days) and cover great distances, with sightings occurring thousands of kilometers away from the main breeding colonies (Aurioles-Gamboa et al. 1999). Guadalupe fur seals are infrequently observed in U.S. waters, but they have been observed at California's Channel Islands.

#### **ESA-Listed Sea Turtles in the Action Area**

The green turtle has a circumglobal distribution, occurring throughout nearshore tropical, subtropical and, to a lesser extent, temperate waters. After emerging from the nest, hatchlings swim to offshore areas and go through a post-hatchling pelagic stage believed to last several years. Adult green turtles exhibit site fidelity and migrate hundreds to thousands of kilometers from nesting beaches to foraging areas. Green turtles spend the majority of their lives on coastal foraging grounds, which include open coastlines and protected bays and lagoons. Figure 11 shows the ranges and boundaries of all green turtle DPSs (Seminoff et al. 2015).



**Figure 11. Map of the 11 Distinct Population Segments of green turtles**

The North Atlantic DPS of green turtle nests primarily outside the action area in Costa Rica and Mexico, but also in Cuba and Florida. Nesting for the green turtle South Atlantic DPS occurs on both sides of the North and South Atlantic Oceans, Ascension Island, and the U.S. Virgin Islands. On the western side of the Atlantic, juveniles and adults can be found on feeding grounds in the Caribbean, which overlaps the action area, and the nearshore waters of Brazil, Uruguay and Argentina, which are not in the action area. The major nesting site for the Central North Pacific DPS of green turtle is at East Island, French Frigate Shoals, in the Northwestern Hawaiian Islands. Central West Pacific DPS green turtle nesting occur in the Federated States of Micronesia and the Marshall Islands. East Pacific DPS nesting occurs outside the action area at major sites in Michoacán, Mexico, and the Galapagos Islands, Ecuador. Smaller nesting sites are found in the Revillagigedo Archipelago, Mexico, and along the Pacific Coast of Costa Rica, Columbia, Ecuador, Guatemala, and Peru. Central South Pacific DPS nesting occurs sporadically throughout the geographic distribution of the population, with isolated locations having relatively low to moderate nesting activity. The Southwest Pacific DPS major nesting sites include the Great Barrier Reef, eastern Torres Strait, and the northern Great Barrier Reef. East Indian-West Pacific DPS nesting sites are in Northern Australia, Indonesia, Malaysia, Peninsular Malaysia and the Philippine Turtle Islands. North Indian DPS major nesting sites are found on the Arabian Peninsula to the Pakistani-Indian border. Major nesting beaches for the Southwest Indian DPS are found in the French Eparses Islands, Mayotte and the outer Seychelles Islands.

The hawksbill turtle has a circumglobal distribution throughout tropical and, to a lesser extent, subtropical waters of the Atlantic, Indian, and Pacific Oceans. In their oceanic phase, juvenile hawksbill turtles can be found in *Sargassum* mats; post-oceanic hawksbills may occupy a range of habitats that include coral reefs or other hard-bottom habitats, seagrass, algal beds, mangrove bays, and creeks (Bjorndal and Bolten 2010; Musick and Limpus 1997).

The Kemp's ridley turtle occurs from the Gulf of America and up along the Atlantic coast of the U.S. (TEWG 2000). The majority of Kemp's ridley turtles nest at coastal Mexican beaches in the

Gulf of Mexico. During spring and summer, juvenile Kemp's ridleys occur in the shallow coastal waters of the northern Gulf from south Texas to north Florida. In the fall, most Kemp's ridleys migrate to deeper or more southern, warmer waters and remain there through the winter (Schmid 1998). As adults, many Kemp's ridley turtles remain in the Gulf, with only occasional occurrence in the Atlantic Ocean (NMFS et al. 2010).

Globally, olive ridley turtles can be found in tropical and subtropical waters in the Atlantic, Indian, and Pacific Oceans. Major nesting beaches are found outside the action area in Nicaragua, Costa Rica, Panama, India, and Suriname. Olive ridleys may forage across ocean basins, primarily in pelagic habitats, on crustaceans, fish, mollusks, and tunicates. The range of the endangered Mexico's Pacific coast breeding population extends as far south as Peru and up to California. Olive ridley turtles of the Mexico's Pacific coast breeding colonies nest outside the action area on arribada beaches at Mismaloya, Ixtapilla, and La Escobilla, Mexico. Solitary nesting takes place all along the Pacific coast of Mexico.

Loggerhead sea turtles are circumglobal and are found in the temperate and tropical regions of the Atlantic, Indian, and Pacific Oceans. The post-hatchling stage is in pelagic waters and juveniles are first in the oceanic zone and later in the neritic zone (i.e., coastal waters). While in their oceanic phase, loggerhead sea turtles undertake long migrations using ocean currents. Adults and subadults occupy nearshore habitat important for foraging and inter-nesting migration. The Northwest Atlantic Ocean DPS of loggerhead sea turtle hatchlings disperse widely, most likely using the Gulf Stream to drift throughout the Atlantic Ocean. Genetic evidence demonstrates that juvenile loggerheads from southern Florida nesting beaches comprise the vast majority (71–88%) of individuals found in foraging grounds throughout the western and eastern Atlantic (Masuda 2010). Northeast Atlantic Ocean DPS loggerheads are found from western Europe to western Africa. Nesting for the Northeast Atlantic Ocean DPS occurs on the islands of the Cape Verde Archipelago, off the coast of western Africa. South Atlantic Ocean DPS loggerheads are found between the eastern coast of South America and the west coast of Africa. Nesting for the South Atlantic Ocean DPS occurs in Brazil, from the state of Rio de Janeiro to Sergipe, with the majority occurring in Bahia. North Pacific Ocean DPS loggerhead sea turtles are found throughout the Pacific Ocean, north of the equator. Their range extends from the West Coast of North America to eastern Asia. Two major juvenile foraging areas have been identified in the North Pacific Basin: Central North Pacific and off Mexico's Baja California Peninsula. Hatchlings from Japanese nesting beaches outside the action area use the North Pacific Subtropical Gyre and the Kurishio Extension to migrate to those foraging grounds (Abecassis et al. 2013; Seminoff et al. 2014). The range of South Pacific Ocean DPS loggerheads extends from eastern Australia to western South America, south of the equator. Nesting for the South Pacific DPS occurs mostly in eastern Australia (e.g., Queensland) and New Caledonia, outside the action area. North Indian Ocean DPS loggerheads are found in the northern Indian Ocean, along the east coast of Africa, the coasts of Yemen, Oman, Iran, Pakistan, India and Southeast Asia. Major nest sites for the North Indian Ocean DPS occurs along the coast of Oman and nearby islands. Masirah Island in Oman is one of the largest loggerhead nesting sites in the world (Baldwin et al. 2003). Southeast Indo-Pacific DPS loggerheads are found in the Indian Ocean, in the waters around western Australia, Indonesia, and Papua New Guinea. Nesting for loggerheads of the Southeast Indo-Pacific Ocean DPS is restricted to western Australia, with three main nesting beaches: Dirk Hartog Island, Murion Island, and North West Cape (Baldwin

et al. 2003). Southwest Indian Ocean DPS loggerheads are found off the coasts of Africa and Madagascar. Nesting is confined to South Africa and Mozambique.

The leatherback sea turtle is unique among sea turtles for its large size and ability to maintain internal warmth (due to thermoregulatory systems), which allows it to range worldwide from tropical into subpolar latitudes. Leatherbacks occur throughout marine waters, from nearshore habitats to oceanic environments (Shoop and Kenney 1992). Leatherback sea turtles migrate long, transoceanic distances between their tropical nesting beaches and the highly productive temperate waters where they forage, primarily on jellyfish and tunicates. Detailed population structure is unknown, but the leatherback distribution is assumed dependent upon nesting beach locations in the Pacific, Atlantic, and Indian Oceans. Movements are largely dependent upon reproductive and feeding cycles and the oceanographic features that concentrate prey, such as frontal systems, eddy features, current boundaries, and coastal retention areas (Benson et al. 2011).

#### **ESA-Listed Invertebrates in the Action Area**

The range of black abalone encompasses coastal waters from Point Arena in Mendocino County, California to Northern Baja California, Bahia Tortugas, and Isla Guadalupe, Mexico. Black abalone are uncommon north of San Francisco (Morris et al. 1980) and south of Punta Eugenia (P. Raimondi, pers. comm. as cited in Butler et al. 2009). Black abalone live on rocky substrates in coastal and offshore island intertidal reefs up to approximately 18 ft (5.5 m) deep (Leighton 2005 as cited in Butler et al. 2009).

White abalone occur between Point Conception in Santa Barbara County, California, to Punta Abreojos, Baja California, Mexico. White abalone occur along the U.S. west coast among offshore islands and banks (particularly Santa Catalina and San Clemente islands) and mainland inshore waters from Point Conception, California south to Punta Abreojos, Baja California, Mexico (Cox 1960; Cox 1962; Bartsch 1940). Individuals have also been found around several Mexican islands including Isla Cedros and Isla Natividad (Guzmán Del Proó 1992). White abalone occupy open low relief rock or boulder habitat surrounded by sand (Tutschulte 1976) at 98.4–213.3 ft (30–65 m) depth (Haaker et al. 1986).

The chambered nautilus is found in coastal reef habitats in the tropical Indo-Pacific including: American Samoa, Australia, Fiji, India, Indonesia, Malaysia, Papua New Guinea, Philippines, Solomon Islands, and Vanuatu. The chambered nautilus is also possibly native to China, Myanmar, Western Samoa, Thailand, and Vietnam. The species is considered a habitat specialist with a patchy distribution, found in association with steep-sloped forereefs and cannot tolerate temperatures above approximately 25°C or depths exceeding around 2,460.6–2,624.7 ft (750–800 m; Miller 2018).

Queen conch can be found in Flower Garden Banks, Florida Keys, Bermuda, and throughout the Caribbean (Horn et al. 2022). Queen conch inhabit a range of habitat types during their life cycle including seagrass beds, sand flats, algal beds, and rubble areas from a couple inches (few centimeters) deep to approximately 98.4 ft (30 m) deep (NMFS 2014).

There are seven species of corals (elkhorn, staghorn, lobed star, boulder star, mountainous star, pillar, and rough cactus) that occur on shallow reefs from south Florida, in the Gulf of America, and throughout the Caribbean. These corals are often found in coastal reefs, but some may be found farther offshore in bank reefs, if it is shallow enough to provide the necessary photic zone habitat.

The range of the proposed sunflower sea star encompasses coastal waters in the northeast Pacific Ocean from the Aleutian Islands to Baja California, Mexico. The sunflower sea star is considered a habitat generalist and most commonly occurs in coastal waters less than 82 ft (25 m), though they can be found to a depth of 1,427 ft (435 m) throughout the known range. They are rare in waters deeper than 394 ft (120 m; Fisher 1928; Gravem et al. 2021; Hemery et al. 2016; Lambert 2000) but have been observed in depths exceeding 1,427.2 ft (435 m) off Oregon and central California (e.g., Keller et al. 2008).

#### **ESA-Listed Fishes in the Action Area**

The Atlantic salmon is an anadromous fish, and the Gulf of Maine DPS is found in watersheds throughout Maine. Gulf of Maine DPS Atlantic salmon migrate nearly 2,500 mi (approximately 4,000 km) in the open ocean to reach feeding areas in the Davis Strait between Labrador and Greenland.

Atlantic sturgeon spawn in freshwater but spend most of their adult life in the marine environment. Atlantic sturgeon occupy ocean waters and associated bays, estuaries, and coastal river systems from Hamilton Inlet, Labrador, Canada, to Cape Canaveral, Florida (ASMFC 2006; Stein et al. 2004). Five DPSs of Atlantic sturgeon are listed under the ESA: Gulf of Maine, New York Bight, Chesapeake Bay, Carolina, and South Atlantic. Juveniles typically spend two to five years in freshwater before eventually becoming coastal residents as subadults (Boreman 1997; Schueller and Peterson 2010; Smith 1985). Atlantic sturgeon exhibit high fidelity to their natal rivers but can undergo extensive mixing in coastal waters (Grunwald et al. 2008; King et al. 2001; Waldman et al. 2002).

Rockfish occupy the eastern Pacific Ocean in waters from California to Alaska. The bocaccio and the yelloweye rockfish are two different long-lived species of rockfish. The listed bocaccio and yelloweye that reside in the Puget Sound/Georgia Basin are their own distinct DPSs.

The Pacific salmon (chinook, coho, chum, and sockeye) and steelhead trout are anadromous fishes and the various ESA-listed DPSs and ESUs (see Table 4) spawn in their natal rivers in Washington, Oregon, and California. Juvenile chinook may reside in freshwater for 12 to 16 months, but some migrate to the ocean as young-of-the-year within eight months of hatching. Chinook salmon spend a few years feeding in the ocean, and sexually mature between the ages of two and seven but are typically three or four years old when they return to spawn, generally in summer or early fall. Coho salmon spend a year in freshwater and then migrate out to the ocean to spend about 1.5 years feeding before returning to spawn, generally in fall or early winter. Sockeye salmon rear in freshwater for one to three years, after which they reach the smolt stage and migrate to the ocean to feed and grow. They typically mature and return to freshwater to spawn in the summer or fall after two to three years at sea, but some return earlier or stay at sea longer, between four and five years. Steelhead trout typically migrate to open marine waters after

spending two years in freshwater. They reside in marine waters for typically two or three years prior to returning to their natal stream as four- or five-year-olds to spawn shortly after river entry from December through April. Young chum salmon (fry) typically migrate directly to estuarine and marine waters soon after they are born and do not reside in freshwater for an extended period. As chum salmon grow larger, they migrate offshore and as they approach maturity, typically between the ages of three and six, they migrate back to spawn in late summer through March.

The eulachon is also an anadromous fish, though smaller than salmonids, that can be found in the continental shelf waters of the eastern Pacific Ocean. Adult and juvenile Southern DPS eulachon typically occupy waters 164–656.2 ft (50–200 m) deep (Gustafson 2016), and up to depths of about 984.3 ft (300 m), from California to the Bering Sea. Southern DPS eulachon return to spawn in rivers south of the Nass River in British Columbia to the Mad River in California.

The giant manta ray occupies tropical, subtropical, and temperate oceanic waters, and productive coastlines where they feed on zooplankton. Giant manta rays are commonly offshore in oceanic waters and may aggregate in shallow waters (less than 32.8 ft [10 m]) to feed. Giant manta rays can dive to depths between 656.2–1,476.4 ft (200–450 m) deep, sometimes exceeding 3,280.8 ft (1,000 m) deep.

The green sturgeon is an anadromous fish that occurs in the nearshore coastal waters to a depth of 360.9 ft (110 m) from Baja California, Mexico to the Bering Sea, Alaska (Erickson and Hightower 2007). Adult Southern DPS green sturgeon enter San Francisco Bay and migrate up the Sacramento River to spawn (Heublein et al. 2009).

The Gulf sturgeon ranges from Lake Pontchartrain in Louisiana east to the Suwannee river system in Florida. Young-of-the-year slowly work their way downstream from where they hatched and arrive in estuaries and river mouths where they will spend their next six years developing (Sulak and Clugston 1999). After six years, Gulf sturgeon will enter the marine environment to forage on benthic (bottom dwelling) invertebrates along the shallow nearshore (6.6–13.1 ft or 2–4 m depth), barrier island passes, and in unknown offshore locations in the Gulf (Carr et al. 1996; Fox et al. 2002; Huff 1975; Ross et al. 2009).

The gulf grouper occurs throughout the subtropical Pacific Ocean from approximately La Jolla, California, to Mazatlán, Sinaloa, Mexico. They are commonly found in the Gulf of California which is not a part of the action area. Young gulf grouper inhabit shallow coastal habitat whereas adults mainly inhabit rocky reefs, seamounts, and kelp beds 16–600 ft (4.9–182.9 m) deep.

The Nassau grouper is distributed from south Florida throughout the Caribbean, and Bermuda. Juveniles inhabit macroalgae, coral clumps, and seagrass beds, and are relatively solitary. As they grow, they occupy progressively deeper areas and offshore reefs, and can be in schools of up to forty individuals. When not spawning, adults are most common in waters less than 328.1 ft (100 m) deep.

The oceanic whitetip shark is a large pelagic shark distributed globally throughout open ocean waters, outer continental shelves, and around oceanic islands, primarily from 10° North to 10°

South, but may be found up to 30° North to 35° South (Young 2016). They occur from the surface to at least 498.7 ft (152 m) deep and display a preference for water temperatures above 20°C.

The scalloped hammerhead shark is found throughout the world and the listed Central and Southwest Atlantic DPS, Eastern Atlantic DPS, and Eastern Pacific DPS live in coastal warm temperate and tropical seas. The species occurs over continental shelves and the shelves surrounding islands, as well as adjacent deep waters, but is seldom found in waters cooler than 22°C (Compagno 1984; Schulze-Haugen et al. 2003). It ranges from the intertidal and surface to depths of up to 1,476.4–1,679.8 ft (450–512 m), with occasional dives to even deeper waters. It has also been documented entering enclosed bays and estuaries. The Central and Southwest Atlantic DPS of scalloped hammerhead shark's range extends from the southeast coast of Florida down to Uruguay, including the Caribbean Sea, but not the Gulf of America. The Eastern Pacific DPS of scalloped hammerhead shark's range extends from the coast of southern California, and south to Ecuador and part of Peru. The Eastern Atlantic DPS ranges along the west coast of Africa, from Morocco to Namibia.

Shortnose sturgeon occur in estuaries, rivers, and the sea along the east coast of North America (Vladykov and Greeley 1963). Their northerly distribution extends from the Saint John River, New Brunswick, Canada, to the Indian River, Florida (Evermann and Bean 1898; Scott and Scott 1988). Some populations rarely leave freshwater while others are known to migrate along the coast between river systems (Altenritter et al. 2015; Dionne et al. 2013; Quattro et al. 2002; Wirgin et al. 2005).

Historically within the U.S., smalltooth sawfish have been captured in estuarine and coastal waters from New York southward through Texas, with the largest number of recorded captures in Florida (NMFS 2010). Recent capture and encounter data suggest that the current distribution is primarily south and southwest Florida from Charlotte Harbor through the Dry Tortugas (Poulakis and Seitz 2004; Seitz and Poulakis 2002). Outside the U.S., smalltooth sawfish have been confirmed in the Bahamas, and sightings off the coasts of Honduras, Belize, Cuba, Guinea Bissau, and Sierra Leone. Juvenile sawfish spend the first two to three years of their lives in shallow waters provided in the lower reaches of rivers, estuaries, and coastal bays (Simpfendorfer et al. 2008; Simpfendorfer et al. 2011). As smalltooth sawfish approach 98.4 inches (250 centimeters), they become less sensitive to salinity changes and begin to move out of protected shallow water embayments and into the shorelines of barrier islands (Poulakis et al. 2011). Adult sawfish typically occur in more open water, marine habitats (Poulakis and Seitz 2004).

#### **Critical Habitat in the Action Area**

This section discusses designated and proposed critical habitat that is either completely encompassed by the action area or is partially within the action area. The following species have critical habitat that overlaps in areas where the only activity that will occur is vessel or aircraft transit:

- Beluga Whale – Cook Inlet DPS
- North Pacific Right Whale

- Rice's Whale (proposed)
- Green Sea Turtle – East Pacific DPS (proposed) and South Atlantic DPS (proposed)
- Hawksbill Sea Turtle
- Atlantic Salmon – Gulf of Maine DPS
- Atlantic Sturgeon – Carolina DPS, Chesapeake Bay DPS, Gulf of Maine DPS, New York Bight DPS, and South Atlantic DPS
- Bocaccio – Puget Sound/Georgia Basin DPS
- Chinook Salmon – California Coastal ESU, Central Valley Spring-Run ESU, Lower Columbia River ESU, Puget Sound ESU, Sacramento River Winter-Run ESU, Snake River Fall-Run ESU, Snake River Spring/Summer Run ESU, Upper Columbia River Spring-Run ESU, and Upper Willamette River ESU
- Chum Salmon – Columbia River ESU and Hood Canal Summer-Run ESU
- Coho Salmon – Central California Coast ESU, Lower Columbia River ESU, Oregon Coast ESU, and Southern Oregon/Northern California Coast ESU
- Eulachon – Southern DPS
- Gulf Sturgeon
- Smalltooth Sawfish – U.S. DPS
- Sockeye Salmon – Ozette Lake ESU and Snake River ESU
- Steelhead – California Central Valley DPS, Central California Coast DPS, Lower Columbia River DPS, Middle Columbia River DPS, Northern California DPS, Puget Sound DPS, Snake River Basin DPS, South-Central California Coast DPS, Southern California DPS, Upper Columbia River DPS, and Upper Willamette River DPS
- Yelloweye Rockfish – Puget Sound/Georgia Basin DPS
- Elkhorn Coral
- Staghorn Coral
- Black Abalone

The following species have critical habitat in the action area where activities beyond just vessel or aircraft transit may occur (e.g., landings with disposal or expended objects, explosive events) and/or are in the vicinity of launch sites:

- Humpback Whale – Central America DPS, Mexico DPS, and Western North Pacific DPS
- Killer Whale – Southern Resident DPS
- North Atlantic Right Whale
- Bearded Seal – Beringia DPS
- Ringed Seal – Arctic subspecies
- Steller Sea Lion – Western DPS
- Green Sea Turtle – North Atlantic DPS (proposed)
- Green Sturgeon – Southern DPS
- Leatherback Sea Turtle
- Loggerhead Sea Turtle – Northwest Atlantic Ocean DPS
- Nassau Grouper
- Boulder Star Coral
- Lobed Star Coral
- Mountainous Star Coral

- Pillar Coral
- Rough Cactus Coral

The majority of these species' critical habitats occur in areas where only vessel transits associated with the action would occur. Below we describe the areas of their critical habitats where additional components of the action may occur.

***Humpback Whale – Western North Pacific DPS, Central America DPS, and Mexico DPS***

NOAA Fisheries designated critical habitat for the endangered Western North Pacific DPS, the endangered Central America DPS, and the threatened Mexico DPS of humpback whales on May 21, 2021 (86 Fed. Reg. 21082; Figure 12, Figure 13, Figure 14). These areas support prey species that are essential to the conservation of the DPSs. The Central America DPS critical habitat overlaps Pacific Zone 4 where barge landings may occur, and a small portion of the critical habitat also overlaps Pacific Zone 2 where objects may be expended or recovered (e.g., stages and fairings). The Mexico DPS critical habitat includes the Central America DPS critical habitat. A unit of the Mexico DPS and Western North Pacific DPS critical habitats occur around the Pacific Spaceport Complex Alaska in Kodiak, Alaska where launches may occur, though has not occurred for some time.




Figure 12. Critical Habitat for Central America DPS humpback whales




Figure 13. Critical Habitat for Mexico DPS humpback whales




Figure 14. Critical Habitat for Western North Pacific DPS humpback whales

#### ***Killer Whale – Southern Resident DPS***

NMFS designated critical habitat for the Southern Resident DPS killer whale in 2006 (71 Fed. Reg. 69054) and revised it in 2021 (86 Fed. Reg. 41668; Figure 15). Coastal waters designated as critical habitat overlap Pacific Zone 4 where barge landings may occur.





Figure 15. Critical Habitat for Southern Resident DPS Killer Whale

***North Atlantic Right Whale***

NMFS designated two units of critical habitat for the North Atlantic right whale. Unit 1 is for foraging habitat in the Gulf of Maine and Georges Bank region (Figure 16), and is in Zone 3c of the action area, which would have no operations with the exception of vessel transits January through July. Unit 2 is for calving, consisting of all marine waters from Cape Fear, North Carolina, southward to approximately 27 NM below Cape Canaveral, Florida (Figure 17). Unit 2 occurs in Zone 3b of the action area, where operations are also restricted to only vessel traffic during January to July. However, Unit 2 is off the coast of CCSFS and, therefore, adjacent the associated launch and landing sites.






Figure 17. North Atlantic Right Whale Critical Habitat Unit 2